

CORRECTION

Open Access

Correction: Cerebral ischemia induces microvascular pro-inflammatory cytokine expression via the MEK/ERK pathway

Aida Maddahi^{1,2}, Lars Edvinsson^{1,2*}

We wish to clarify in results page 4, line 22 of our study [1]; that the information on infarct volume and neurology score after MCAO was taken from our previous study [2].

Author details

¹Division of Vascular Research, BMC, Lund University, Lund, Sweden.

²Department of Internal Medicine, Institute of Clinical Sciences, Lund University, Lund, Sweden.

Received: 2 February 2011 Accepted: 21 February 2011

Published: 21 February 2011

References

1. Maddahi A, Edvinsson L: Cerebral ischemia induces microvascular pro-inflammatory cytokine expression via the MEK/ERK pathway. *J Neuroinflammation* 7:14.
2. Maddahi A, Edvinsson L: Enhanced expressions of microvascular smooth muscle receptors after focal cerebral ischemia occur via the MAPK MEK/ERK pathway. *BMC Neurosci* 2008, 9:85.

doi:10.1186/1742-2094-8-18

Cite this article as: Maddahi and Edvinsson: Correction: Cerebral ischemia induces microvascular pro-inflammatory cytokine expression via the MEK/ERK pathway. *Journal of Neuroinflammation* 2011 8:18.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



* Correspondence: lars.edvinsson@med.lu.se

¹Division of Vascular Research, BMC, Lund University, Lund, Sweden
Full list of author information is available at the end of the article