

RESEARCH

Open Access

# Changes in TNF $\alpha$ , NF $\kappa$ B and MnSOD protein in the vestibular nuclei after unilateral vestibular deafferentation

Martine Liberge\*, Christine Manrique, Laurence Bernard-Demanze, Michel Lacour

## Abstract

**Background:** Unilateral vestibular deafferentation results in strong microglial and astroglial activation in the vestibular nuclei (VN) that could be due to an inflammatory response. This study was aimed at determining if markers of inflammation are upregulated in the VN after chemical unilateral labyrinthectomy (UL) in the rat, and if the inflammatory response, if any, induces the expression of neuroprotective factors that could promote the plasticity mechanisms involved in the vestibular compensation process. The expressions of inflammatory and neuroprotective factors after chemical or mechanical UL were also compared to verify that the inflammatory response was not due to the toxicity of sodium arsenite.

**Methods:** Immunohistological investigations combined the labeling of tumor necrosis factor  $\alpha$  (TNF $\alpha$ ), as a marker of the VN inflammatory response, and of nuclear transcription factor  $\kappa$ B (NF $\kappa$ B) and manganese superoxide dismutase (MnSOD), as markers of neuroprotection that could be expressed in the VN because of inflammation. Immunoreactivity (Ir) of the VN cells was quantified in the VN complex of rats. Behavioral investigations were performed to assess the functional recovery process, including both static (support surface) and dynamic (air-righting and landing reflexes) postural tests.

**Results:** Chemical UL (arsenite transtympanic injection) induced a significant increase in the number of TNF $\alpha$ -Ir cells in the medial and inferior VN on both sides. These changes were detectable as early as 4 h after vestibular lesion, persisted at 1 day, and regained nearly normal values at 3 days. The early increase in TNF $\alpha$  expression was followed by a slightly delayed upregulation of NF $\kappa$ B 8 h after chemical UL, peaking at 1 day, and regaining control values 3 days later. By contrast, upregulation of MnSOD was more strongly delayed (1 day), with a peak at 3 days, and a return to control values at 15 days. Similar changes of TNF $\alpha$ , NF $\kappa$ B, and MnSOD expression were found in rats submitted to mechanical UL. Behavioral observations showed strong posturo-locomotor deficits early after chemical UL (1 day) and a complete functional recovery 6 weeks later.

**Conclusions:** Our results suggest that the upregulation of inflammatory and neuroprotective factors after vestibular deafferentation in the VN may constitute a favorable neuronal environment for the vestibular compensation process.

## Background

Vestibular compensation, i.e. functional recovery after a unilateral lesion of the peripheral vestibular system, is a good model to study the cellular and molecular mechanisms involved in the neuroplasticity of the adult

central nervous system. Unilateral loss of vestibular inputs to the brainstem vestibular nuclei (VN) is caused by lesioning the peripheral sensory receptors (chemical or mechanical unilateral labyrinthectomy, UL) or unilaterally transecting the vestibular nerve (UVN). These procedures produce a complex vestibular syndrome made of static and dynamic signs. The static signs include postural deficits (increased support surface, head tilt) and ocular motor deficits (spontaneous nystagmus) that are compensated within a few days or weeks,

\* Correspondence: martine.liberge@univ-provence.fr  
Université Aix-Marseille, UMR 6149 Université de Provence/CNRS, Neurobiologie Intégrative et Adaptative, Pôle 3C, Comportement, Cerveau, Cognition, Centre de St Charles, Case B, 3 Place Victor Hugo, 13331 Marseille Cedex 3, France

depending on the species and on the type of vestibular deafferentation. Recent findings in the cat indicated that vestibular deficits resulting from a functional deafferentation are compensated faster (2 weeks after mechanical UL) than those induced by an anatomical deafferentation (about 2 months after UVN) [1]. On the contrary, the dynamic signs (vestibulo-ocular reflex asymmetry, postural instability and equilibrium) are compensated much less completely or over a longer time [2,3].

Several hypotheses have been proposed to explain the vestibular compensation process [2,4]. At least part of the behavioral recovery process has been attributed to plasticity of neuronal and non-neuronal elements in the deafferented VN. In line, recent findings have demonstrated reactive neurogenesis in the adult cat [5,6] and a strong microglial and astroglial reaction [5,7-9] after unilateral deafferentation in the deafferented VN. Campos-Torres et al. [9] suggested that the glial reaction may be due to modifications of the environment of VN neurons and most probably to the central inflammatory tissue reaction consecutive to UL. Despite this inflammatory reaction, the central vestibular neurons did not show any sign of apoptosis after the permanent loss of their peripheral inputs [8]. Glial reaction in the vestibular nuclei may be a beneficial mechanism. It could both protect the injured tissue from long-lasting inflammation and promote the survival of deafferented vestibular neurons [9] or of newly generated neurons [6]. Finally, the inflammatory reaction could help in the vestibular compensation process.

Upon insult or infection, the brain inflammatory response generally involves the production of proinflammatory cytokines such as tumor necrosis factor  $\alpha$  (TNF $\alpha$ ), interleukins (IL1 $\beta$ , IL-6...), interferon gamma (IFN- $\gamma$ ), and various reactive oxygen and nitrogen species [10]. To date, the cellular inflammatory mediators produced by central vestibular glial cells and/or neurons after peripheral vestibular loss have not been characterized.

Among the inflammatory mediators, TNF $\alpha$  is a prominent effector of the immune response in the brain. It is constitutively expressed by astrocytes, microglia, and neurons in the normal brain [11] and upregulated following various manifestations of brain injury, including neurodegenerative disorders (Alzheimer's and Parkinson's diseases, multiple sclerosis), brain trauma, and ischemic injury (for review, see [12]). It has been suggested TNF $\alpha$  may thus have a pathogenic role in these disorders. Nevertheless, TNF $\alpha$  has been shown to exert both neurotoxic and neuroprotective effects in experimental models. TNF $\alpha$  can mediate cell death [13,14] but, conversely, TNF $\alpha$  is thought to enhance neuroprotective processes in models of acute neurodegeneration (ischemia, excitotoxicity, axotomy) and chronic (Alzheimer's disease) neurodegeneration [15-20].

Increased levels of TNF $\alpha$  protect cells against damage from cytotoxic insults in part via the induction of cytoprotective genes, including Manganese Superoxide Dismutase (MnSOD) [21-23]. The antioxidant enzyme MnSOD is critical for neuronal survival in various paradigms of stress-induced brain injury [23-26]. Xu et al. [27] demonstrated that the presence of an intronic nuclear factor  $\kappa$ B (NF $\kappa$ B) site in the MnSOD gene was essential for the induction of MnSOD by TNF $\alpha$ . NF $\kappa$ B is a nuclear transcription factor that has a protective role in many tissues in response to stress, inflammation, and immunity. NF $\kappa$ B protects neurons from toxic insults and promotes neuronal growth and synaptic plasticity [24,28-30]. Widera et al. [31] demonstrated that NF $\kappa$ B plays a crucial role in the proliferation of neural stem cells induced by TNF $\alpha$ .

The present study aimed to determine whether a chemical UL (transtympanic injection of sodium arsenite) inflicts an inflammatory response in the vestibular nuclei. This reaction could be expressed by an increased production of proinflammatory cytokines such as TNF $\alpha$  and could induce a compensatory change in the expression of the neuroprotective factors NF $\kappa$ B and MnSOD during the course of the vestibular recovery. This was achieved using immunohistochemistry, notably by quantifying modulation of TNF $\alpha$ , NF $\kappa$ B, and MnSOD labeling in the four main ipsilateral and contralateral VN (medial, superior, lateral, and inferior). The immunohistochemical data were recorded in groups of rats killed 4 h, 8 h, 1 day, 3 days, 7 days, and 15 days after chemical UL. In addition, functional recovery after chemical UL was tested in another group of rats.

TNF $\alpha$ , NF $\kappa$ B, and MnSOD labeling was also quantified in the ipsilateral and contralateral medial (MVN) and inferior (IVN) nuclei in groups of rats submitted to mechanical UL and sacrificed 8 h, 1, and 7 days after the lesion. Effects of mechanical and chemical UL were compared to ensure that arsenite-induced changes in the expression of inflammatory and neuroprotective factors were not due to the toxicity of arsenite on brainstem structures.

## Methods

### Animals care and housing

The experiments were performed on male adult Long Evans rats (300-350 g) obtained from R. Janvier (Le Genest-St-Isle, France). Animals were maintained on a 12 h light/dark cycle at a constant temperature (22°C) with ad libitum access to food and water. The experiments were performed in accordance with the requirements of the French "Ministère de l'Agriculture et de la Pêche" Décret no. 87-848, October 19, 1987 and to the European Communities Council Directive of November 24, 1986 (86/609/EEC). Every attempt was made to

minimize both the number and suffering of animals used in these experiments.

### Vestibular lesions

Chemical UL was produced by a unilateral transtympanic injection of the toxin, sodium arsanilate. Sodium arsanilate (Sigma, St Louis, MO) was dissolved in 0.3 M sodium carbonate at the concentration of 300 mg/ml and titrated to pH 7.5.

The rats were anesthetized with an intramuscular injection of a solution of ketamine (Ketamine 1000, Virbac, Carros, France, 62.5 mg/kg) and medetomidine (Domitor, Orion Pharma, Espoo, Finland, 0.4 mg/kg). Guided through a dissecting microscope, a 26-gauge needle of injection was advanced through the tympanic membrane, and 100  $\mu$ l of sodium arsanilate was infused in the left middle-ear. After the injection needle was withdrawn, the external ear canal was packed with sterile hemostatic sponge to prevent leakage of the toxin away from the injection site. When the surgery was completed, the effects of medetomidine were reversed by a subcutaneous injection of the specific antagonist atipamezole (0.33 mg/kg) (Pfizer, Paris, France). Using this protocol, the effects of anesthesia were rapidly reversed and, a few minutes later, the UL rats were able to stand and thereafter to behave nearly normally.

The mechanical UL was performed using a retroauricular approach. The entire length of the ventral edge of the left meatus was destroyed. The tympanic membrane, malleus, incus, and the stapes were removed to expose the pterygopalatine. This artery was coagulated using an electrocoagulator and the oval window was opened. Finally, the vestibule was drilled and aspirated using a suction tube. A postoperative i.m. Terramycin (Pfizer, Paris, France) injection was given to prevent infection.

### Histological preparation

Thirty rats were subjected to a chemical UL. They received unilateral transtympanic injection of sodium arsanilate on the left side and were killed at six recovery times after vestibular lesion: 4 h ( $n = 5$ ), 8 h ( $n = 5$ ), 1 day ( $n = 5$ ), 3 days ( $n = 5$ ), 7 days ( $n = 5$ ), 15 days ( $n = 5$ ). Sham-operated rats ( $n = 24$ , four rats per group) received a unilateral transtympanic injection of 0.9% sterile saline and were killed at the same recovery time as lesioned animals.

Six rats were submitted to mechanical UL on the left side. Based on data recorded in the chemical UL rats, they were killed at three recovery times: 8 h ( $n = 2$ ), 1 day ( $n = 2$ ), 7 days ( $n = 2$ ).

Rats were deeply anesthetized with sodium pentobarbital (100 mg/kg IP) and perfused intracardially with 100 ml of 0.9% saline followed by 400 ml of 4% paraformaldehyde (Sigma, St Louis, MO) in 0.1 M phosphate

buffer (PB), pH 7.4. The brains were rapidly removed and postfixed in a 4% paraformaldehyde solution for 24 h, cryoprotected by successive transfers into increasing concentrations (10%, 20%, and 30%) of sucrose solution in 0.1 M PB for 72 h at 4°C. Brains were then rapidly frozen on powdered dry ice (CO<sub>2</sub> gas) and kept at -80°C until use. They were sectioned coronally at 30  $\mu$ m on a cryostat, and the sections were stored into stock solution (0.1 M PB (pH 7.4), 30% (v/v) glycerol and 30% (v/v) ethylene glycol) at -20°C until immunohistochemistry experiments.

### Immunohistochemistry (IHC)

Free-floating sections were rinsed three times with 0.1 M phosphate buffer saline (PBS) for 5 min, preincubated in a PBS 0.1 M solution containing 0.25% Triton X-100 and 10% bovine serum albumin (BSA) for 1 h at room temperature. Thereafter, sections were incubated under continuous agitation 24 h at 4°C in either a goat polyclonal TNF $\alpha$  antibody (1:200), a rabbit polyclonal NF $\kappa$ B antibody (1:200), or a rabbit monoclonal MnSOD antibody (1:200), diluted in a PBS 0.1 M solution containing 0.25% Triton X-100 and 2% BSA. The antibodies for TNF $\alpha$  and NF $\kappa$ B were obtained from Santa Cruz Biotechnology (Santa Cruz, CA), anti-MnSOD from Epitomic (Burlingame, CA).

On the following day, sections were washed 3  $\times$  5 min in PBS 0.1 M, BSA 2% and 3  $\times$  5 min in PBS 0.1 M, BSA 5%. Then, they were incubated in 1:200 biotinylated secondary antibody diluted in PBS 0.1 M, 2% BSA for 1 h at room temperature. Sections then underwent another round of washing with PBS 0.1 M, BSA 2% (3  $\times$  5 min) and with PBS 0.1 M, BSA 5% (3  $\times$  5 min) and were incubated in biotinylated horseradish peroxidase avidin (Vector Laboratories, Burlingame, CA) for 1 h at room temperature. Finally, for visualization, sections were reacted using 0.2 mg/ml diaminobenzidine and 0.03% hydrogen peroxide in PBS 0.1 M. After being washed with PB, the sections were mounted on microscope slides and allowed to dry overnight. The following day, sections were dehydrated in ascending series of alcohol concentrations (70%, 80%, 90%, and 100%), cleared with xylene, and coverslipped with DPX mounting media.

Immunolabeling for TNF $\alpha$ , NF $\kappa$ B, and MnSOD was examined 4 h, 8 h, 1, 3, 7 and 15 days after chemical UL, and 8 h, 1 and 7 days after mechanical UL. The specificity of the immunolabeling was validated by omission of the primary antibody. No labeling was observed in these control conditions.

### Data quantification of TNF $\alpha$ , NF $\kappa$ B, and MnSOD immunoreactivity in the vestibular nuclei

The vestibular nuclei were identified according to Paxinos and Watson's stereotaxic atlas of the rat [32].

The four VN (medial, superior, lateral, and inferior; hereinafter MVN, SVN, LVN, and IVN respectively) were easily differentiated without counterstaining. Data quantification in the MVN, SVN, LVN, and IVN was done in serial sections collected from the rostral (-9.96 mm) to the caudal (-11.76 mm) part of the brainstem. Stained cells were counted in the MVN (27 sections per rat), SVN (24 sections), LVN (15 sections), and IVN (24 sections). The variable number of sections depended on the rostrocaudal extent of the nuclei [5]. The immunoreactivity quantification was performed on both sides.

The number of cells expressing TNF $\alpha$ , NF $\kappa$ B, or MnSOD was quantified via computer-assisted image analysis using a DMLB microscope (Leica Microsystems, Wetzlar, Germany) equipped with a DXM 1200 Nikon high-resolution digital camera (1024-1024 pixels; Nikon, Tokyo, Japan) interfaced to a PC computer employing image software for capturing and processing the images (Lucia G, Nikon, Champigny-sur-Marne, France). Quantification of the labeled cells was performed via a grey-level method by adjusting a threshold brightness value, and only cells labeled with a grey value above this threshold were taken into account. Reproducibility was assessed by comparing the same data analyzed independently by two researchers blind to the animal groups. Specific labeling was quantified in each section as the number of labeled cells and was automatically computed and expressed as the mean ( $\pm$  SEM) for each side, each rat, and each subgroup of rats [5]. To eliminate quantification problems due to potentially asymmetric slides, data were collected from symmetrical slides only. The absence of structural asymmetry in the slides was verified on slides stained with cresyl violet.

### Behavioral investigations

Behavioral investigations were performed in another group of rats ( $n = 6$ ) to assess static and dynamic postural deficits after arsanilate lesion and the time course of the functional recovery.

Static postural deficits and recovery were evaluated by measuring the surface delimited by the four legs of the rat standing erect at rest, without walking. Support surface is a very sensitive parameter used for evaluating vestibular lesion-induced static posture deficit and recovery. To quantify the support surface, the rats were placed in a box equipped with a transparent bottom and filmed from under this box. A scale drawn on the bottom served to take measurements of the four paws location. Four repeated measurements were taken for each rat before chemical UL, and at regular intervals during the course of recovery until complete recovery (60 days). The support surface was measured using an image analysis system (Canvas 9TM, Deneba Software, Miami, FL). An average was calculated for each postlesion time. Data recorded after vestibular lesions were compared to the

mean prelesion values using individual references, that is, each animal acted as its own control.

Dynamic postural deficits and recovery were evaluated in the same group of rats ( $n = 6$ ) using two dynamic reflex tests, the air-righting reflex and the landing reflex. The air-righting reflex was used to test the rat's ability to right itself in the air, while the landing reflex was used to test the response of the rat to a vertical linear acceleration. The tests were considered to evaluate the level of the vestibular deficit and of the recovery process following the vestibular lesion.

During the air-righting test, the rats were held at 50 cm above a cushion, in supine position. The experimenter removed his/her hands as quickly and simultaneously as possible. In a normal rat, just after the drop, the vestibular system detects a change in linear acceleration that triggers the head turning, which in turn induces the sequence of body repositioning. After vestibular lesion, the righting response is totally abolished and it recovers with time through the vestibular compensation process. A score of "0" was assigned when the air-righting reflex was complete, "2" when the animal failed to return itself and fell on its back, and "1" when it showed a partial reaction only.

During the landing reflex, the rats were held by the base of the tail and lowered toward the ground. Animals with intact vestibular function flex their neck and extend their forelimbs as they approach the surface, whereas rats with vestibular lesion at an uncompensated stage show the right response only when their forepaws or vibrissae touch the ground. A score of "0" was assigned when the animals responded properly, and "2" if ground contact was required.

The scores for the air-righting reflex and the landing reflex were summed, providing a global score of dynamic vestibular function and recovery. A zero score indicated a normal vestibular function as observed before the vestibular lesion, or a complete recovery after UL. Scores ranging from 1 to 4 pointed to vestibular dysfunction and more or less complete vestibular compensation. Three repeated measurements were taken for statokinetic tests for each rat before chemical UL and at each postoperative time. A mean was calculated for each postlesion time and compared to the mean prelesion values. As for the support surface evaluation, each animal acted as its own control.

For some experimental animals, double-blind quantification was performed for the dynamic reflex tests, which did not show any dependence on the experimenter when he/she was trained to perform the tests.

### Statistical analysis

TNF $\alpha$ , NF $\kappa$ B, and MnSOD immunolabeling in the VN observed after a chemical UL was analyzed using

repeated-measures analysis of variance (ANOVAs) with lesion (sham vs arsanilate-lesioned rats) and postoperative recovery period (4, 8 h, and 1, 3, 7, 15 days) as between-factors, and with the nuclei (MVN, SVN, LVN and IVN) as within factors. When a significant difference was revealed, Student's t-tests were used for further two by two comparisons between sham and lesioned animals at the six postoperative times. The levels of significance were set at  $p < 0.01$  (TNF $\alpha$  and NF $\kappa$ B) and  $p < 0.0083$  (MnSOD) according to the number of multiple comparisons (ie, Bonferroni's principle).

TNF $\alpha$ , NF $\kappa$ B, and MnSOD immunolabeling in the MVN and IVN after mechanical and chemical UL was analyzed using repeated-measures ANOVA with type of lesion (mechanical vs chemical UL) and postoperative time (8 h, 1 and 7 days) as between-factors, and with the vestibular nuclei (MVN and IVN) as within factors. Unpaired t-tests were used for comparisons between mechanical and chemical UL animals at the three postoperative times. The levels of significance were set at  $p < 0.016$  (TNF $\alpha$ , NF $\kappa$ B, and MnSOD) according to the Bonferroni's principle.

Support surface values at each postoperative time were compared with preoperative surface values for each rat by a paired Student's t-test. The global score of the rats in the air-righting and landing reflex tests recorded at each postoperative time was compared to the preoperative performance of each rat by a paired Student's t-test. The levels of significance were set at  $p < 0.0038$  (support surface and global score) according to the number of multiple comparisons (Bonferroni's principle).

## Results

### 1. Immunohistochemical data

#### *Chemical UL-induced change in TNF $\alpha$ immunoreactivity in the vestibular nuclei*

Repeated-measures ANOVA of the cell counts of TNF $\alpha$ -positive cells demonstrated that the lesion ( $p < 0.0001$ ), the structure (MVN, SVN, LVN and IVN) ( $p < 0.0001$ ), and the postoperative time (4 h, 8 h, 1, 3, 7 and 15 days) ( $p < 0.05$ ) constituted the main fixed effects providing the sources of variation among animals. This was also corroborated by the significant interaction between these variables ( $p < 0.0001$ ). In contrast, no significant differences were found between the ipsilateral and contralateral VN for both the sham-operated and the arsanilate-lesioned rats. Consequently, the data from both sides were pooled and averaged for both groups of rats.

Figure 1A illustrates the TNF $\alpha$ -Ir changes in the MVN and the IVN of two representative rats: one sham-operated rat and one arsanilate-lesioned rat killed 8 h after the vestibular lesion. Chemical UL resulted in an increase in TNF $\alpha$  expression in the both VN, as seen

by the increased number of TNF $\alpha$ -Ir cells. This change in TNF $\alpha$  expression was also observed in the SVN, but not in the LVN.

Quantitative analysis of TNF $\alpha$ -like immunolabeling in the MVN and the IVN is shown in Figure 1B. A significantly increased number of labeled cells was first detected 4 h following the lesion, but the highest significant numbers of TNF $\alpha$ -positive cells were encountered in the MVN ( $288.6 \pm 11.9$ ,  $p < 0.0001$ ) and IVN ( $75.9 \pm 1.2$ ,  $p < 0.001$ ) at the 8 h survival period. One day after the lesion, TNF $\alpha$ -like immunoreactivity persisted significantly ( $249.5 \pm 25.4$  ( $p < 0.01$ ) and  $69.4 \pm 4.5$  ( $p < 0.01$ ) for the MVN and IVN, respectively).

The mean number of TNF $\alpha$ -Ir cells regained normal values three days after the lesion ( $173.8 \pm 16.4$  and  $50.9 \pm 5.6$ , in the MVN and IVN respectively).

#### *Chemical UL-induced change in NF $\kappa$ B immunoreactivity in the vestibular nuclei*

Repeated-ANOVA revealed significant effects of the lesion ( $p < 0.05$ ) and an interaction between the lesion and the postoperative time ( $p < 0.05$ ). Data obtained in the sham-operated groups at the different postoperative times were pooled and considered as control values.


Figure 2 shows microphotographs of NF $\kappa$ B-Ir sections taken for the ipsilateral MVN and IVN from one representative sham-operated rat and from one rat killed 1 day after the lesion. As observed with TNF $\alpha$ , the number of NF $\kappa$ B-Ir cells in both VN increased substantially after arsanilate UL.

Quantitative analysis of the data is shown in Figure 3 for the MVN and the IVN, displayed here as representative nuclei of the VN complex. The average number of NF $\kappa$ B-Ir cells in the MVN and IVN was  $177.1 \pm 11.5$  and  $68.5 \pm 5.9$ , respectively, in the controls. A significant bilateral increase was first observed 8 h postlesion in the MVN ( $258.2 \pm 30.6$ ,  $p < 0.01$ ) and remained unchanged at 1 day ( $267.3 \pm 37.9$  for MVN,  $p < 0.01$ ). Three days after arsanilate lesion, the mean number of NF $\kappa$ B-Ir cells tend to return to control values in the MVN ( $244.6 \pm 32.0$ ). The same time-course was observed in the IVN, even though it remained just below the level of significance ( $p = 0.03$  and  $p = 0.02$  at 8 h and 1 day respectively).



#### *Chemical UL-induced change in MnSOD immunoreactivity in the vestibular nuclei*

Repeated-ANOVA revealed significant effects of the lesion ( $p < 0.001$ ), the structure ( $p < 0.0001$ ), and the postoperative time ( $p < 0.01$ ); it also showed that the interaction between these factors ( $p < 0.05$ ) was significant. As for NF $\kappa$ B quantification, data obtained in the sham-operated group at the different postoperative times were pooled and expressed as control values.

Arsanilate UL resulted in an increase in MnSOD expression in the bilateral VN. An illustration of this



**Figure 1** Chemical UL-induced changes of TNF $\alpha$ -Ir cells in the MVN and IVN. (A) Illustration of the typical labeling in a representative sham rat and in an experimental animal subjected to chemical UL (arsanilate transtympanic injection) observed 8 h after the lesion. Note that chemical UL induces an increased number of TNF $\alpha$ -Ir cells. Scale bar: 100  $\mu$ m. (B) Quantitative analysis of the effects of chemical UL on TNF $\alpha$ -Ir cells in the MVN and IVN. Data are mean values ( $\pm$  SEM) of the number of TNF $\alpha$ -Ir recorded in the sham rats (black histograms) and in the UL rats (grey histograms) observed at 4, 8 h, 1, 3, 7, and 15 days after the lesion. The data from both sides were pooled for sham and experimental animals. According to Bonferroni's principle,  $P < 0.01$  was considered significant. \*  $P < 0.01$ , \*\*  $P < 0.001$ , \*\*\*  $P < 0.0001$  versus sham values (MVN: medial vestibular nucleus; IVN: inferior vestibular nucleus).



effect in the MVN and IVN, 3 days postoperatively is shown in Figure 4A. This change of MnSOD expression was also observed in the SVN but not in the LVN.

Figure 4B illustrates the quantitative analysis of these data in the MVN and the IVN. Compared to controls, which exhibited a mean number of MnSOD-Ir cells of  $86.6 \pm 3.7$  for the MVN and  $37.9 \pm 0.8$  for IVN, the 1 day postlesion group displayed an increase in MnSOD-labeled cells with a mean number of  $127.6 \pm 6.1$  and  $52.8 \pm 0.6$  for the MVN and IVN respectively ( $p < 0.00083$ ). The number of MnSOD-Ir cells continued to increase at 3 days postlesion ( $170.2 \pm 13.4$  and  $60.6 \pm 2.4$  for the MVN ( $p < 0.00083$ ) and for the IVN ( $p < 0.00083$ ) respectively), and remained unchanged in the MVN at 7 days ( $167.9 \pm 17.3$ ,  $p < 0.00083$ ). Fifteen days after arsanilate UL, normal values were recorded in the MVN ( $96.4 \pm 7.0$ ) and the IVN ( $42.6 \pm 2.9$ ).

#### **Upregulation of TNF $\alpha$ , NF $\kappa$ B, and MnSOD expression after chemical and mechanical UL**

Repeated ANOVA of the cell counts of TNF $\alpha$ , NF $\kappa$ B, or MnSOD-positive cells revealed no significant effect of the type of lesion (chemical vs mechanical) and no significant interaction between the lesion and the postoperative time for all the markers. No significant differences were found between the ipsilateral and contralateral MVN and IVN of mechanical UL rats at the different postoperative times. Consequently, data from both sides were pooled and averaged for each group of mechanical UL rats.

Quantitative analysis of TNF $\alpha$ , NF $\kappa$ B, or MnSOD immunolabeling in the MVN and IVN after mechanical and chemical UL is shown in Figure 5. Whatever the postoperative times, the mean values in the mechanical UL rats did not differ significantly from those of the chemical UL rats (Figure 5).

For the TNF $\alpha$ -positive cells, the mean numbers recorded in the 8 h postoperative group were  $288.6 \pm 11.9$  and  $218.5 \pm 6.7$  in the MVN for the chemical and mechanical UL rats, respectively (NS). A similar upregulation of TNF $\alpha$ -Ir cells was observed with both types of lesions at the 1 day survival period in the MVN (Figure 5A). Similar changes were found in the IVN when considering the chemical and mechanical groups of rats (Figure 5A), and when considering the NF $\kappa$ B-positive cells (Figure 5B) as well as the MnSOD-positive cells (Figure 5C) in both the MVN and IVN. Only one significant difference ( $p < 0.016$ ) was seen for the MnSOD-positive cells in the IVN at the 1 day survival period, in favor of the mechanical UL group. In addition, return towards normal values was similar for the three markers considered in the present study in both the chemical and mechanical UL rats, indicating a similar time-course of upregulation in both models of unilateral vestibular deafferentation.

## **2. Behavioral investigations**

### **Behavioral observations**

Unilateral inner ear injection of arsanilate induced the typical behavioral signs of vestibular deafferentation. All arsanilate-UL animals exhibited characteristic oculomotor nystagmus with slow phase directed toward the lesioned side and fast phase beating toward the contralateral side. The postural and locomotor symptoms consisted of rolling and circling behaviors, head tilt and falls toward the lesioned side, and postural asymmetry with strong enhancement of the support surface.

### **Recovery of static posture function**

Recovery of static posture function after arsanilate-UL was evaluated by measuring the support surface delimited by the four legs of the rat standing erect at rest. Figure 6 shows the time-course of posture function recovery after vestibular lesion. The support surface was rather small in normal rats ( $20.2 \pm 1.0 \text{ cm}^2$ ) and greatly increased in the days following the vestibular lesion. Maximal support surface increase ( $50.1 \pm 1.1$ ,  $p < 0.00038$ ) was observed 1 day postlesion, resulting from a strong hypotonia of the limb extensors ipsilateral to the lesion and strong hypertonia of the contralateral extensors muscles. The support surface decreased progressively during the following days but remained superior to preoperative values on the 8<sup>th</sup> postoperative day ( $38.6 \pm 1.5$ ,  $p < 0.00038$ ). At 30 days postlesion, the support surface remained significantly larger ( $27.0 \pm 1.2$ ,  $p < 0.0038$ ). The complete compensation of this postural parameter required 36 days.

### **Recovery of dynamic posture function**


Before vestibular lesion, all animals had the ability to right themselves when they were held in a supine position and dropped in the air. The animals also succeeded in flexing their neck and extending their forelimbs when they were lowered toward the ground. The air-righting and the landing reflexes were normal in these controls and the global score was zero.

After arsanilate-UL, rats failed to succeed in either test. Figure 7 shows the mean time-course of recovery. One day after the lesion, the mean global score of the UL rats was  $2.3 \pm 0.2$  ( $p < 0.00038$ ), indicating that animals failed in performing correct body air-righting reflex and landing reflex. From day 2 to 8, the mean global score remained significantly increased ( $2.9 \pm 0.1$ ,  $p < 0.00038$ ) on day 8. Then, the UL rats' deficits gradually decreased over time and animals fully recovered both air-righting reflex and landing reflex as correctly as normal rats 42 days after UL only.


## **Discussion**

The present study shows first that a chemical UL induces the typical postural and locomotor symptoms seen after all types of vestibular deafferentation and a


**A**




**B**




**Figure 4 Chemical UL-induced changes of MnSOD-Ir cells in the MVN and IVN.** (A) Illustration of a typical labeling in a representative sham-operated rat and in a chemical UL animal examined 3 days after the lesion, showing an increased number of MnSOD-Ir cells in both VN. Scale bar: 100 µm. (B) Quantitative analysis of the effects of chemical UL on MnSOD-Ir cells in the MVN and IVN. Data are mean values ( $\pm$  SEM) of the number of MnSOD-Ir cells in the MVN and IVN for rats that had chemical UL and observed at 4, 8 h, 1, 3, 7, and 15 days after the lesion. Data of all the postoperative times were pooled in sham animals, and mean value is shown as a horizontal line and standard error of the mean ( $\pm$  SEM) as two dotted lines. According to Bonferroni's principle,  $P < 0.0083$  was considered significant. \*  $P < 0.0083$ , \*\*  $P < 0.00083$  versus sham values.



**Figure 5 Comparative TNF $\alpha$ , NF $\kappa$ B and MnSOD immunostaining in the MVN and the IVN after a chemical and a mechanical UL.** Quantitative analysis of the effects of chemical and mechanical UL on TNF $\alpha$ -Ir cells (A), NF $\kappa$ B-Ir cells (B) and MnSOD-Ir (C) in the MVN and IVN. Data are mean values ( $\pm$  SEM) of the number of positive cells recorded in the chemical UL rats (grey histograms) and in the mechanical UL rats (black histograms) observed at 8 h, 1 and 7 days after the lesions. The data from both sides were pooled for all groups of animals. According to Bonferroni's principle,  $P < 0.016$  was considered significant. \*  $P < 0.016$  mechanical versus chemical values.



**Figure 6 Recovery of static posture function after chemical UL.** The support surface (expressed in cm<sup>2</sup>) was evaluated for each rat (n = 4 measurements per rat) at regular time intervals until complete recovery. Each point represents the mean value ( $\pm$  SEM) for each postoperative time calculated in a group of chemical UL rats (n = 6). The mean of support surface preoperative values is shown as a horizontal line and standard error of the mean ( $\pm$  SEM) as two dotted lines. Note the strong increase in surface in the days following chemical UL and the complete recovery after 36 days. According to Bonferroni's principle,  $P < 0.0038$  was considered significant. \*  $P < 0.0038$ , \*\*  $P < 0.00038$  versus preoperative values.



**Figure 7 Recovery of dynamic posture function after chemical UL.** Each point represents the global score of the rats to the air-righting and landing reflex tests at each postoperative time and was compared to preoperative values. A zero score indicated a normal vestibular function, scores ranging from 1 to 4 pointed to vestibular dysfunction and less complete vestibular compensation. Before chemical UL (pre-op), the mean global score was zero. After the lesion, rats failed in performing correct body air-righting and landing reflexes. Recovery was complete 42 days after UL. According to Bonferroni's principle,  $P < 0.0038$  was considered significant. \*  $P < 0.0038$ , \*\*  $P < 0.00038$ , versus preoperative values.

recovery of both static and dynamic posture functions requiring 5-6 weeks. Second, the present findings demonstrate that chemical UL induces a rapid inflammatory reaction in the VN, as shown by the significantly increased expression of the proinflammatory cytokine TNF $\alpha$ . This TNF $\alpha$  upregulation is followed by an increased expression of the antioxidant enzyme MnSOD, associated with the upregulation of the nuclear transcription factor NF $\kappa$ B. Similar changes of TNF $\alpha$ , NF $\kappa$ B, and MnSOD expression were found in rats submitted to mechanical UL. The changes in the expression of neuroprotective factors could constitute a favorable neuronal environment for the vestibular compensation process.

### 1. Behavioral recovery after chemical UL

Unilateral transtympanic injection of sodium arsenite in the rat induces classical postural deficits (ipsilateral head tilt, increased support surface with ipsilateral limb flexion and contralateral limb extension) and locomotor (ipsilateral falls, rolling and circling) deficits. However, the complete recovery of these postural functions is much longer than that reported in this species after mechanical UL [9,33]. Indeed, head tilt and postural asymmetry disappeared 3-4 days after mechanical UL [9], while 6 weeks were required in the present study. These differences in the time constant of the recovery process were already reported by Kim et al. [34], who showed head deviation peaking 2 h and 3 days after mechanical UL and chemical UL, respectively. In addition, these authors reported a head posture significantly tilted during at least one week after chemical UL since longer recovery delays were not tested in their study. Interestingly, the unilateral section of the vestibular nerve (UVN) in the cat model also induced a delayed behavioral recovery (6 weeks) compared to mechanical UL (2 weeks) [1]. Such differences in the recovery profiles can be accounted for by the nature of the VN deafferentation. There is a total, sudden, anatomical and functional deafferentation after UVN whereas mechanical UL leads to a partial and progressive anatomical deafferentation [1,35]. And it was recently reported that the plasticity mechanisms involved in the recovery process also differ with the type of vestibular deafferentation [36].

A first explanatory hypothesis is that chemical UL is a model of VN deafferentation that is closer to the UVN model. Indeed, intratympanic injection of sodium arsenite has been shown to cause permanent loss of vestibular function as a result both of degeneration of the vestibular sensory epithelium [37,38] and of the vestibular nerve [39]. This degeneration of the vestibular nerve would explain why the recovery process is longer after chemical UL than after mechanical UL. However, recent

findings in the adult rat submitted to chemical UL, similar to our protocol, clearly showed that the Scarpa's ganglion neurons, the vestibular nerve, and the brainstem structures did not exhibit any degeneration after intratympanic injection of sodium arsenite, from 3 days to 1 month after injection (Chabbert, personal communication). The transtympanic injection targets on epithelial vestibular sense organ (and on cochlear epithelium) and toxicity of arsenite were similar to that of glutamate excito-toxicity [40]. In this case, the differences between mechanical and chemical UL could be in the postlesion discharge in the vestibular nerve. No electrophysiological inputs to the vestibular nuclei are seen just after mechanical UL while abnormal electrophysiological action potentials could occur in the vestibular nerve after arsenite lesion, due to axonal terminations still connected to hair cells. This hypothesis is based on the observations of heterotopic electrical discharges in the hair cells after glutamate excito-toxicity [40]. If confirmed, this hypothesis could explain the longer recovery time-course after chemical UL than mechanical UL in rodents.

### 2. Inflammatory and neuroprotective responses in the VN after unilateral vestibular deafferentation

A massive microglial and astroglial reaction was observed in the bilateral VN after mechanical UL in the rat [8,9]. Moreover, axotomy or removal of the peripheral sensory hair cells failed to cause any neuronal cell death in the VN [8]. Not only the second-order vestibular neurons do not degenerate, but these neuronal cells recover a near normal resting discharge one week after mechanical UL in the guinea-pig [41]. The reactive gliosis observed in the VN could be therefore a beneficial mechanism protecting the injured nervous tissue. As suggested by Campos-Torres et al. [9] and described in other nerve injury paradigms [42,43], pro- and anti-inflammatory cytokines produced by activated glial cells could promote the survival of deafferented vestibular neurons and therefore may contribute to the vestibular compensation process.

#### *Chemical UL-induced upregulation of TNF $\alpha$ in the VN*

Our findings clearly showed a significant increase in the proinflammatory cytokine TNF $\alpha$  in the bilateral VN, detectable as early as 4 h after UL and persisting until 3 days. TNF $\alpha$  is expressed by neurons, in addition to glial cells, in the normal brain, and TNF $\alpha$  receptors are present on all neural cell types, rendering the cells responsive to TNF $\alpha$  signaling [11]. TNF $\alpha$  is upregulated in several neurodegenerative disorders (cerebral malaria, AIDS dementia, Alzheimer's disease) and has been primarily considered to promote the pathologies [44-46]. Besides this potential neurotoxic effect, numerous studies have revealed an unexpected role of TNF $\alpha$  in CNS

repair. It protects the brain against ischemia or nitric oxide-induced injury [18,20], and glutamate [17,19,47] or  $\beta$  amyloid neurotoxicity [15,23], in part by induction of antioxidant pathway [15]. Neuronal damage after focal cerebral ischemia or administration of kainic acid is increased in mice lacking TNF $\alpha$  receptor [48]. After optic nerve axotomy, TNF $\alpha$  was found to prevent secondary death of the retinal ganglion cells [16]. Such a neuroprotective effect of TNF $\alpha$  could promote the survival of both the present VN neurons (no cell death) and the newly generated neurons found in the deafferented VN after UVN in the cat [5,6]. The wide variety of responses to TNF $\alpha$ , including apoptosis as well as cell survival and cell proliferation, can be in part explained by signaling pathways mediated through different TNF receptors (TNFR). TNFR1 contains an intracellular "death domain" associated with the detrimental effects of TNF $\alpha$  in models of acute neurodegeneration [13,49,50] while TNFR2 would contribute to cell survival, growth, and proliferation [13,17,51-53].

#### **Chemical UL-induced upregulation of NF $\kappa$ B in the VN**

NF $\kappa$ B is a transcription factor activated after brain ischemia [54] and spinal cord injury [55,56] and in neurodegenerative processes [57]. However, like TNF $\alpha$ , NF $\kappa$ B also has a neuroprotective role. Preactivation of NF $\kappa$ B by ceramide or by low amounts of TNF $\alpha$  protects neurons from glutamate ototoxicity or oxidative insults [58]. NF $\kappa$ B activation mediates the anti-death action of TNF $\alpha$  in hippocampal neurons exposed to hypoxia or  $\beta$  amyloid toxicity [15,24,28]. Such a TNF $\alpha$ -induced NF $\kappa$ B activation could play a similar neuroprotective action after chemical UL. Indeed, our results showed that NF $\kappa$ B was upregulated in the bilateral VN shortly after the upregulation of TNF $\alpha$ . In cortical neurons exposed to glutamate toxicity, the signaling pathways involve TNFR1 and TNFR2 [17]. While activation of the TNFR1 pathway leads to a transient NF $\kappa$ B upregulation (30 min), TNFR2 signaling promotes persistent and long lasting NF $\kappa$ B activation (up to 24 h) [17]. These authors suggested that the duration of NF $\kappa$ B activation was critical to achieve significant tissue protection. Since NF $\kappa$ B was upregulated in the VN up to 20 h after onset of TNF $\alpha$  upregulation, we hypothesize that TNF $\alpha$  could act through TNFR2 to activate neuroprotective signaling pathways involved in the recovery process after chemical UL.

#### **Chemical UL-induced upregulation of MnSOD in the VN**

A delayed and long lasting upregulation of MnSOD was shown also in the bilateral VN after chemical UL. These changes were detectable 1 day after arsanilate injection and persisted up to 7 days. MnSOD is regulated transcriptionally and induced by inflammatory cytokines, including TNF $\alpha$ , IL1, and IFN $\gamma$  [59-61]. Yune et al. [22] demonstrated that exogenous TNF $\alpha$  induces MnSOD expression in the uninjured spinal cord, a process that is

inhibited after treatment with a TNF $\alpha$  antibody in injured tissue.

Inflammatory cytokines like TNF $\alpha$  generate reactive oxygen species (ROS), and the mitochondrial respiratory chain is the main source of TNF $\alpha$ -induced ROS [62]. The cells contain a variety of antioxidant enzymes, such as mitochondrial MnSOD, which defend against free radicals damage. MnSOD protects cells against oxidative stress by rapidly dismutating superoxide radicals (O $_2$ ) to hydrogen peroxide (H $_2$ O $_2$ ) [63]. Hydrogen peroxide can be further detoxified into water and oxygen by catalase or glutathione peroxidase [64]. Accumulating data indicate that MnSOD is critical for neuronal survival against various stress-induced brain injuries [25,65], and that increased levels of TNF $\alpha$  in response to cytotoxic insults protect neurons in part via the induction of cytoprotective genes, including MnSOD [22-24,26]. MnSOD protection has also been demonstrated in non-neuronal cells [21,27,66].

Little is known, however, on the molecular mechanisms that govern transcription of the MnSOD gene. NF $\kappa$ B has been proposed as one of the three transcription factors (with AP-1 and C/EBP- $\beta$ ) regulating TNF $\alpha$ -induced MnSOD expression in different cell types [21-23,62,66]. It is likely that MnSOD induction, found in the present study in VN cells after chemical UL, is mediated through activation of NF $\kappa$ B. However, NF $\kappa$ B and MnSOD expressions showed different kinetics, with NF $\kappa$ B peaking at 1 day postlesion while MnSOD expression peaked 3 days postlesion, as previously reported after spinal cord injury [22]. This suggests that other transcription factors could be involved. Before being activated, NF $\kappa$ B resides in the cytoplasm in an inactive form consisting of three subunits: p50 and p65 (constituting the factor dimer p50-p65) and the I $\kappa$ B inhibitory subunit. When I $\kappa$ B is bound to p50-p65, NF $\kappa$ B is inactive. Signals activating NF $\kappa$ B (TNF $\alpha$ : see [67]) cause dissociation of this complex, releasing p50-p65 that then translocates to the nucleus and binds to specific  $\kappa$ B DNA sequences in the promoter region of a variety of NF $\kappa$ B-responsive genes, such as MnSOD gene. Xu et al. [27] found that induction of MnSOD by TNF $\alpha$  required the presence of an intronic nuclear factor  $\kappa$ B (NF $\kappa$ B) site in the MnSOD gene. And protection of neuronal cells from  $\beta$ -amyloid peptide toxicity by TNF $\alpha$ -induced MnSOD activation resulted from an enhanced binding of the dimer p50-p65 to the promoter/enhancer regions of the MnSOD gene [23]. The upregulation of the neuroprotective MnSOD protein observed in our chemical UL model could be due to activation of such a sequence.

#### **Effects of chemical and mechanical UL on TNF $\alpha$ , NF $\kappa$ B, and MnSOD upregulation in the rat**

Comparison of chemical and mechanical UL data strongly suggests that the inflammatory response

observed after chemical UL is not due to a toxic effect per se of sodium arsenite on brainstem structures. Since the amount as well the kinetics of the upregulation of the three markers did not differ for the mechanical UL group compared to the chemical UL group, this suggests that the cellular response to vestibular deafferentation is similar in both models.

These results are supported by Campos-Torres and de Kazak's personal observations, reported in the discussion of the paper by Campos-Torres et al., 2005 [9]. The authors studied cytokine expression within the vestibular nuclei at different times after mechanical labyrinthectomy in the rat. They reported that the real time light cycler RT-PCR studies showed a significant upregulation of TNF $\alpha$  and interleukin (IL1, IL-6) within the vestibular nuclei beginning at 6 h after the lesion.

#### **Changes of TNF $\alpha$ , NF $\kappa$ B and MnSOD protein in the bilateral VN**

Why the changes in all the markers (TNF $\alpha$ , NF $\kappa$ B, and MnSOD) can be seen bilaterally, given that the lesion was unilateral, can be explained with respect to the neural connectivity between the VN on both sides, and to their cell content. Neuroanatomical studies have shown that the VN neurons are under the control of both excitatory inputs from the ipsilateral labyrinth and inhibitory inputs from the contralateral labyrinth via direct projections (commissural system) and indirect ones (cerebellar cortex loops) [68]. It was also shown that the VN contains different subpopulations of neurons, including excitatory second order vestibular cells (tonic and kinetic type I neurons), inhibitory second order vestibular cells, excitatory interneurons, and inhibitory interneurons (type II neurons), at least for the MVN [69]. Vestibular deafferentation can therefore lead to changes in different subpopulations of cells in the VN on the intact and lesioned sides. Dysinhibitory and/or dysfacilitatory processes can activate type I cells on the intact side, as evidenced electrophysiologically [70], whereas facilitatory processes can activate both excitatory and inhibitory interneurons on the lesioned side. Such mechanisms can also explain the early bilateral increase in c-Fos mRNA and Fos protein levels observed in the MVN, SVN, and IVN after hemilabyrinthectomy in the rat [71]. Indeed, those authors observed that the changes were asymmetric in the MVN, being most prominent in the dorsal part of the contralateral MVN (where second order vestibular neurons are located) and in the ventral part of the ipsilateral nucleus (where commissural neurons acting on the MVN of the intact side are located). Many other works also pointed out bilateral changes in the cat model for c-Fos [72], choline acetyltransferase-Ir [73], and GABA-Ir [[74], see 36 for a review].

## **Conclusions**

Our study indicates that behavioral recovery after chemical UL in the rat is similar to that previously reported in the cat model after UVN. We provide cellular evidence that this type of vestibular deafferentation, as well as mechanical UL, induces a strong inflammatory reaction in the VN, as shown by the increased expression of the proinflammatory cytokine TNF $\alpha$ . The compensatory responses to this stress, that is, upregulation of the neuroprotective factors NF $\kappa$ B and MnSOD, very likely constitute a favorable environment 1) to prevent existing VN cells from apoptosis, and 2) to promote the plasticity mechanisms involved in the recovery process, including cell survival and cell differentiation of newly generated cells. Indeed, the role of TNF $\alpha$  and NF $\kappa$ B in cell proliferation and cell differentiation has been strengthened by recent reports in both neonatal [75] and adult [31,76] brain structures. The present findings strongly suggest that proinflammatory cytokines and induced neuroprotective factors play a role in the recovery of vestibular functions.

#### **List of abbreviations used**

ANOVA: analysis of variance; BSA: bovine serum albumin; IVN: inferior vestibular nucleus; LVN: lateral vestibular nucleus; MVN: medial vestibular nucleus; NF $\kappa$ B: nuclear transcription factor  $\kappa$ B; MnSOD: manganese superoxide dismutase; SVN: superior vestibular nucleus; TNF $\alpha$ : tumor necrosis factor alpha; PB: phosphate buffer; PBS: phosphate buffer saline; UL: unilateral labyrinthectomy; UVN: unilateral vestibular neurectomy; VN: vestibular nuclei.

#### **Acknowledgements**

This study was supported by grants from the Ministère de l'enseignement supérieur et de la recherche and CNRS (UMR Université de Provence, CNRS No. 6149). The authors thank Valerie Gilbert and Catherine Marra for taking care of the animals.

#### **Authors' contributions**

ML(\*) is the main investigator of this work in charge of the study design, analysis and interpretation of results, and writing. CM and LBD both participated in immunohistological investigations and analysis of the data. ML(\*\*) participated mainly in the writing of the report. All authors read and approved the final manuscript.

#### **Competing interests**

The authors declare that they have no competing interests.

Received: 5 July 2010 Accepted: 9 December 2010

Published: 9 December 2010

#### **References**

1. Lacour M, Dutheil S, Tighilet B, Lopez C, Borel L: **Tell Me Your Vestibular Deficit, and I'll Tell You How You'll Compensate.** *Ann NY Acad Sci* 2009, **1164**:268-278.
2. Dieringer N: **Vestibular compensation: neural plasticity and its relations to functional recovery after labyrinthine lesion in frogs and other vertebrates.** *Prog Neurobiol* 1995, **46**:97-129.
3. Darlington CL, Smith PF: **Molecular mechanisms of recovery from vestibular damage in mammals: recent advances.** *Prog Neurobiol* 2000, **62**:13-25.
4. Smith PF, Curthoys IS: **Mechanisms of recovery following unilateral labyrinthectomy: a review.** *Brain Res Rev* 1989, **14**:155-180.

5. Tighilet B, Brezun JM, Gustave Dit Duflo S, Gaubert C, Lacour M: **New neurons in the vestibular nuclei complex after unilateral vestibular neurectomy in the adult cat.** *Eur J Neurosci* 2007, **25**:47-58.
6. Dutheil S, Brezun JM, Leonard J, Lacour M, Tighilet B: **Neurogenesis and astrogenesis contribution to recovery of vestibular functions in the adult cat following unilateral vestibular neurectomy: cellular and behavioral evidence.** *Neuroscience* 2009, **164**:1444-1456.
7. Li H, Godfrey DA, Rubin AM: **Astrocyte reaction in the rat vestibular nuclei after unilateral removal of Scarpa's ganglion.** *Ann Otol Rhinol Laryngol* 1999, **108**:181-188.
8. Campos-Torres A, Vidal PP, De Waele C: **Evidence for a microglial reaction within the vestibular and cochlear nuclei following inner ear lesion in the rat.** *Neuroscience* 1999, **92**:1475-1490.
9. Campos-Torres A, Touret M, Vidal PP, Barnum S, De Waele C: **The differential response of astrocytes within the vestibular and cochlear nuclei following unilateral labyrinthectomy or vestibular afferent activity blockade by transtympanic tetrodotoxin injection in the rat.** *Neuroscience* 2005, **130**:853-865.
10. Rock RB, Gekker G, Hu S, Sheng WS, Cheeran M, Lokensgard JR, Peterson PK: **Role of microglia in central nervous system infections.** *Clin Microbiol Rev* 2004, **17**:942-964.
11. Vitkovic L, Bockaert J, Jacque C: **"Inflammatory" cytokines: neuromodulators in normal brain?** *J Neurochem* 2000, **74**:457-471.
12. Shoshani E, Ginis I, Hallenbeck JM: **Dual role of tumor necrosis factor in brain injury.** *Cytokine Growth Factor Rev* 1999, **10**:119-130.
13. Fontaine V, Mohand-Said S, Hanoteau N, Fuchs C, Pfizenmaier K, Eisel U: **Neurodegenerative and neuroprotective effects of tumor necrosis factor (TNF) in retinal ischemia: Opposite roles of TNF receptor 1 and TNF receptor 2.** *J Neurosci* 2002, **22**:RC216.
14. Kenchappa P, Yadav A, Singh G, Nandana S, Banerjee K: **Rescue of TNF alpha-inhibited neuronal cells by IGF-1 involves Akt and c-Jun N-terminal kinases.** *J Neurosci Res* 2004, **76**:466-474.
15. Barger SW, Horster D, Furukawa K, Goodman Y, Kriegstein J, Mattson MP: **Tumor necrosis factors  $\alpha$  and  $\beta$  protect neurons against amyloid  $\beta$ -peptide toxicity: Evidence for involvement of a  $\kappa$ B-binding factor and attenuation of peroxide and  $\text{Ca}^{2+}$  accumulation.** *Proc Natl Acad Sci* 1995, **92**:9328-9332.
16. Diem R, Meyer R, Weishaupt JH, Bähr M: **Reduction of potassium currents and phosphatidylinositol 3-kinase-dependent Akt phosphorylation by tumor necrosis factor- $\alpha$  rescues axotomized retinal ganglion cells from retrograde cell death in vivo.** *J Neurosci* 2001, **21**:2058-2066.
17. Marchetti L, Klein M, Schlett K, Pfizenmaier K, Eisel ULM: **Tumor necrosis factor (TNF)-mediated neuroprotection against glutamate induced excitotoxicity is enhanced by N-methyl-D-aspartate receptor activation. Essential role of a TNF receptor 2-mediated phosphatidylinositol 3-kinase-dependent NF  $\kappa$ B pathway.** *J Biol Chem* 2004, **279**:32869-32881.
18. Turrin NP, Rivest S: **Tumor necrosis factor  $\alpha$  but not interleukin  $1\beta$  mediates neuroprotection in response to acute nitric oxide excitotoxicity.** *J Neurosci* 2006, **2**:143-151.
19. Dolga AM, Granic I, Blank T, Knaus HG, Spiess J, Luiten PG, Eisel UL, Nijholt IM: **TNF- $\alpha$  mediates neuroprotection against glutamate-induced excitotoxicity via NF- $\kappa$ B-dependent up-regulation of  $\text{K}_{\text{Ca}}2.2$  channels.** *J Neurochem* 2008, **107**:1158-1167.
20. Lambertsen KL, Clausen BH, Babcock AA, Gregersen R, Fenger C, Nielsen HH, Haugaard LS, Wrenfeldt M, Nielsen M, Dagnaes-Hansen F, Bluethmann H, Faergeman NJ, Meldgaard M, Deierborg T, Finsen B: **Microglia protect neurons against ischemia by synthesis of tumor necrosis factor.** *J Neurosci* 2009, **29**:1319-1330.
21. Jones PL, Ping D, Boss JM: **Tumor necrosis factor alpha and interleukin-1 beta regulate the murine manganese superoxide dismutase gene through a complex intronic enhancer involving C/EBP-beta and NF-kappaB.** *Mol Cell Biol* 1997, **17**:6970-6981.
22. Yune TY, Lee SM, Kim SJ, Park HK, Oh YJ, Kim YC, Markelonis GJ, Oh TH: **Manganese superoxide dismutase induced by TNF- $\alpha$  is regulated transcriptionally by NF- $\kappa$ B after spinal cord injury in rats.** *J Neurotrauma* 2004, **21**:1778-1795.
23. Sompol P, Xu Y, Ittarat W, Daosukho C, St Clair D: **NF- $\kappa$ B-associated MnSOD induction protects against  $\beta$ -Amyloid-induced neuronal apoptosis.** *J Mol Neurosci* 2006, **29**:279-288.
24. Mattson MP, Goodman Y, Luo H, Fu W, Furukawa K: **Activation of NF- $\kappa$ B protects hippocampal neurons against oxidative stress-induced apoptosis: evidence for induction of manganese superoxide dismutase and suppression of peroxynitrite production and protein tyrosine nitration.** *J Neurosci Res* 1997, **49**:681-697.
25. Keller JN, Kindy MS, Holtzman FW, St Clair DK, Yen HC, Germeyer A, Steiner SM, Bruce-Keller AJ, Hutchins JB, Mattson MP: **Mitochondrial MnSOD prevents neural apoptosis and reduces ischemic brain injury: suppression of peroxynitrite production, lipid peroxidation and mitochondrial dysfunction.** *J Neurosci* 1998, **18**:687-697.
26. Bruce-Keller AJ, Geddes JW, Knapp PE, Keller JN, Holtzman FW, Steiner SM, Mattson MP: **Anti-death properties of TNF against metabolic poisoning: mitochondrial stabilization by MnSOD.** *J Neuroimmunol* 1999, **93**:53-71.
27. Xu Y, Kinningham KK, Devalaraja MN, Yeh CC, Majima H, Kasarskis EJ, St Clair DK: **An intronic NF- $\kappa$ B element is essential for induction of the human manganese superoxide dismutase gene by tumor necrosis factor- $\alpha$  and interleukin-1 beta.** *DNA Cell Biol* 1999, **18**:709-722.
28. Kaltschmidt B, Uherek M, Wellmann H, Volk B, Kaltschmidt C: **Inhibition of NF- $\kappa$ B potentiates amyloid  $\beta$ -mediated neuronal apoptosis.** *Proc Natl Acad Sci* 1999, **96**:9409-9414.
29. Albeni BC, Mattson MP: **Evidence for the involvement of TNF and NF- $\kappa$ B in hippocampal synaptic plasticity.** *Synapse* 2000, **35**:151-159.
30. Foehr ED, Lin X, O'Mahony A, Gelezianus R, Bradshaw RA, Greene WC: **NF- $\kappa$ B signaling promotes both cell survival and neurite process formation in nerve growth factor-stimulated PC12 cells.** *J Neurosci* 2000, **20**:7556-7563.
31. Widera D, Mikenberg I, Elvers M, Kaltschmidt C, Kaltschmidt B: **Tumor necrosis factor alpha triggers proliferation of adult neural stem cells via IKK/NF- $\kappa$ B signaling.** *BMC Neurosci* 2006, **7**:64-82.
32. Paxinos G, Watson C: *The Rat Brain in Stereotaxic Coordinates, third edition* Academic Press Inc, New York; 1997.
33. Sirkin DW, Precht W, Courjon JH: **Initial, rapid phase of recovery from unilateral vestibular lesion in rat not dependent on survival of central portion of vestibular nerve.** *Brain Res* 1984, **302**:245-256.
34. Kim KMS, Kim JH, Jin YZ, Kry D, Park BR: **Temporal changes of cFos-like protein expression in medial vestibular nuclei following arsanilate-induced unilateral labyrinthectomy in rats.** *Neurosci Lett* 2002, **319**:9-12.
35. Schuknecht HF: **Behavior of the vestibular nerve following labyrinthectomy.** *Ann Otol Rhinol Laryngol Suppl* 1982, **97**:16-32.
36. Lacour M, Tighilet B: **Plastic events in the vestibular nuclei during vestibular compensation: the brain orchestration of a "deafferentation" code.** *Restor Neurol Neurosci* 2010, **28**:19-35.
37. Annik M, Wersäll J: **Experimentally (atoxyl) induced ampullar degeneration and damage to the maculae utriculi.** *Acta Otolaryngol* 1977, **83**:429-440.
38. Kaufman GD, Anderson JH, Beitz AJ: **Fos-defined activity in rat brainstem following centripetal acceleration.** *J Neurosci* 1992, **12**:4489-4500.
39. Chen YC, Pellis SM, Sirkin DW, Potegal M, Teitelbaum P: **Backfall: Labyrinthine and Non-Labyrinthine Components.** *Physiol Behav* 1986, **37**:805-814.
40. Brugeaud A, Travo C, Demêmes D, Lenoir M, Llorens J, Puel JL, Chabbert C: **Control of Hair Excitability by the Vestibular Primary Sensory Neurons.** *J Neurosci* 2007, **27**:3503-3511.
41. Ris L, De Waele C, Serafin M, Vidal PP, Godaux E: **Neuronal activity in the ipsilateral vestibular nucleus following unilateral labyrinthectomy in the alert guinea pig.** *J Neurophysiol* 1995, **74**:2087-2099.
42. Raivich G, Bohatschek M, Kloss CU, Werner A, Jones LL, Kreutzberg GW: **Neuroglial activation repertoire in the injured brain: graded response, molecular mechanisms and cues to physiological function.** *Brain Res Brain Res Rev* 1999, **30**:77-105.
43. Raivich G, Jones LL, Werner A, Bluthmann H, Doetschmann T, Kreutzberg GW: **Molecular signals for glial activation: pro and anti-inflammatory cytokines in the injured brain.** *Acta Neurochir Suppl (Wien)* 1999, **73**:21-30.
44. Tyor WR, Wesselingh SL, Griffin JW, McArthur JC, Griffin DE: **Unifying hypothesis for the pathogenesis of HIV-associated dementia complex, vacuolar myelopathy, and sensory neuropathy.** *J Acquired Immune Defic Syndr Hum Retroviro* 1995, **9**:379-388.
45. Lou J, Lucas R, Grau GE: **Pathogenesis of cerebral malaria: recent experimental data and possible applications for humans.** *Clin Microbiol Rev* 2001, **14**:810-820.
46. Perry RT, Collins JS, Wiener H, Acton R, Go RC: **The role of TNF and its receptors in Alzheimer's disease.** *Neurobiol Aging* 2001, **22**:873-883.

47. Shinpo K, Kikuchi S, Moriwaka F, Tashiro K: **Protective effects of the TNF-ceramide pathway against glutamate neurotoxicity on cultured mesencephalic neurons.** *Brain Res* 1999, **819**:170-173.
48. Bruce AJ, Boling W, Kindy MS, Peschon J, Kraemer PJ, Carpenter MK, Holtzman FW, Mattson MP: **Altered neuronal and microglial responses to excitotoxic an ischemic brain injury in mice lacking TNF receptors.** *Nat Med* 1996, **2**:788-794.
49. Kassiotis G, Kollias G: **Uncoupling the proinflammatory from the immunosuppressive properties of tumor necrosis factor (TNF) at the p55 TNF receptor level. Implications for pathogenesis and therapy of autoimmune demyelination.** *J Exp Med* 2001, **193**:427-434.
50. Yang L, Lindholm K, Konishi Y, Li R, Shen Y: **Target depletion of distinct tumor necrosis factor receptor subtypes reveals hippocampal neuron death and survival through different signal transduction pathways.** *J Neurosci* 2002, **22**:3025-3032.
51. Shen Y, Li R, Shiosaki K: **Inhibition of p75 tumor necrosis factor receptor by antisense oligonucleotides increases hypoxic injury and beta-amyloid toxicity in human neuronal cell line.** *J Biol Chem* 1997, **272**:3550-3553.
52. Arnett HA, Mason J, Marino M, Suzuki K, Matsushima GK, Ting JPY: **TNF $\alpha$  promotes proliferation of oligodendrocyte progenitors and remyelination.** *Nat Neurosci* 2001, **4**:1116-1122.
53. Dolga AM, Nijholt IM, Ostroveanu A, Ten Bosch Q, Luiten PG, Eisel ULM: **Lovastatin induces neuroprotection through tumor necrosis factor receptor 2 signaling pathways.** *J Alzheimers Disease* 2008, **13**:111-122.
54. Clemens JA, Stephenson DT, Smalstig EB, Dixon EP, Little SP: **Global ischemia activates nuclear factor-kB in forebrain neurons of rats.** *Stroke* 1997, **28**:1073-1080.
55. Xu J, Fan G, Chen S, Wu Y, Xu XM, Hsu CY: **Methylprednisolone inhibition of TNF- $\alpha$  expression and NF- $\kappa$ B activation after spinal cord injury in rats.** *Brain Res Mol Brain Res* 1998, **59**:135-142.
56. Kim GM, Xu J, Xu J, Song SK, Yan P, Ku G, Xu XM, Hsu CY: **Tumor necrosis factor receptor deletion reduces nuclear factor-kappaB activation, cellular inhibitor of apoptosis protein 2 expression, and functional recovery after traumatic spinal cord injury.** *J Neurosci* 2001, **21**:6617-6625.
57. Kaltschmidt B, Uherek M, Volk B, Baeuerle PA, Kaltschmidt C: **Transcription factor NF- $\kappa$ B is activated in primary neurons by amyloid beta peptides and in neurons surrounding early plaques from patients with Alzheimer disease.** *Proc Natl Acad Sci* 1997, **94**:2642-2647.
58. Goodman Y, Mattson MP: **Ceramide protects hippocampal neurons against excitotoxic and oxidative insults, and amyloid  $\beta$ -peptide toxicity.** *J Neurochem* 1996, **66**:869-872.
59. Visner GA, Dougall WC, Wilson JM, Burr IA, Nick HS: **Regulation of manganese superoxide dismutase by lipopolysaccharide, interleukin-1, and tumor necrosis factor. Role in the acute inflammatory response.** *J Biol Chem* 1990, **265**:2856-2864.
60. Harris CA, Derbin KS, Hunte-McDonough B, Krauss MR, Chen KT, Smith DM, Epstein LB: **Manganese superoxide dismutase is induced by IFN- $\gamma$  in multiple cell types. Synergistic induction by IFN- $\gamma$  and tumor necrosis factor or IL-1.** *J Immunol* 1991, **147**:149-154.
61. Karube-Harada A, Sugino N, Kashida S, Takiguchi S, Takayama H, Yamagata Y, Nakamura Y, Kato H: **Induction of manganese superoxide dismutase by tumor necrosis factor- $\alpha$  in human endometrial stromal cells.** *Mol Hum Reprod* 2001, **7**:1065-1072.
62. Delhalle S, Deregowski V, Benoit V, Merville MP, Bours V: **NF- $\kappa$ B-dependent MnSOD expression protects adenocarcinoma cells from TNF- $\alpha$ -induced apoptosis.** *Oncogene* 2002, **21**:3917-3924.
63. Weisiger RA, Fridovich I: **Mitochondrial superoxide dismutase. Site of synthesis and intramitochondrial localization.** *J Biol Chem* 1973, **248**:4793-4796.
64. Michiels C, Raes M, Toussaint O, Remacle J: **Importance of Se-glutathione peroxidase, catalase, and Cu/Zn-SOD for cell survival against oxidative stress.** *Free Radic Biol Med* 1994, **17**:235-248.
65. Gonzalez-Zulueta M, Ensz LM, Mukhina G, Lebovitz RM, Zwacka RM, Engelhardt JF, Oberley LW, Dawson VL, Dawson TM: **Manganese superoxide dismutase protects nNOS neurons from NMDA and nitric oxide-mediated neurotoxicity.** *J Neurosci* 1998, **18**:2040-2055.
66. Manna SK, Zhang HJ, Yan T, Oberley LW, Aggarwal BB: **Overexpression of manganese superoxide dismutase suppresses tumor necrosis factor-induced apoptosis and activation of nuclear transcription factor-kappaB and activated protein-1.** *J Biol Chem* 1998, **273**:13245-13254.
67. Chen ZJ, Hagler J, Palombella VJ, Melandri F, Scherer D, Ballard D, Maniatis T: **Signal-induced site specific phosphorylation targets I kappa B alpha to the ubiquitin proteasome pathway.** *Genes Dev* 1995, **9**:1586-1597.
68. Pompeiano O, Mergner T, Corvaja N: **Commissural, perihypoglossal and reticular afferent projections to the vestibular nuclei in the cat. An experimental anatomical study with the method of the retrograde transport of horseradish peroxidase.** *Archs Ital Biol* 1978, **116**:13-172.
69. Wilson VJ, Wylie RM, Marco LA: **Synaptic inputs to cells in the medial vestibular nucleus.** *J Neurophysiol* 1967, **31**:176-185.
70. Shimazu H, Precht W: **Inhibition of central vestibular neurons from the contralateral labyrinth and its mediating pathway.** *J Neurophysiol* 1966, **29**:467-492.
71. Cirelli C, Pompeiano M, D'Ascanio P, Arrighi P, Pompeiano O: **c-fos expression in the rat brain after unilateral labyrinthectomy and its relation to the uncompensated and compensated stages.** *Neuroscience* 1996, **70**:515-546.
72. Gustave-Dit-Duflo S, Gestreau C, Tighilet B, Lacour M: **Fos expression in the cat brainstem after unilateral vestibular neurectomy.** *Brain Res* 1999, **824**:1-17.
73. Tighilet B, Lacour M: **Distribution of choline acetyltransferase immunoreactivity in the vestibular nuclei of normal and unilateral vestibular neurectomized cats.** *Eur J Neurosci* 1998, **10**:3115-3126.
74. Tighilet B, Lacour M: **Gamma amino butyric acid (GABA) immunoreactivity in the vestibular nuclei of normal and unilateral vestibular neurectomized cats.** *Eur J Neurosci* 2001, **13**:2255-2267.
75. Bernardino L, Agasse F, Silva B, Ferreira R, Grade S, Malva JO: **Tumor necrosis factor- $\alpha$  modulates survival, proliferation, and neuronal differentiation in neonatal subventricular zone cell cultures.** *Stem cells* 2008, **26**:2361-2371.
76. Denis-Donini S, Dellarole A, Crociara P, Francese MT, Bortolotto V, Quadrato G, Canonico PL, Orsetti M, Ghi P, Memo M, Bonini SA, Ferrari-Toninelli G, Grilli M: **Impaired adult neurogenesis associated with short-term memory defects in NF- $\kappa$ B p50-deficient mice.** *J Neurosci* 2008, **28**:3911-3919.

doi:10.1186/1742-2094-7-91

**Cite this article as:** Liberge et al.: Changes in TNF $\alpha$ , NF $\kappa$ B and MnSOD protein in the vestibular nuclei after unilateral vestibular deafferentation. *Journal of Neuroinflammation* 2010 **7**:91.

**Submit your next manuscript to BioMed Central and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
www.biomedcentral.com/submit

