

RESEARCH

Open Access



Simplified regimen of combined low-dose rituximab for autoimmune encephalitis with neuronal surface antibodies

Ying Du^{1†}, Chao Zhao^{1†}, Juntong Liu^{1†}, Chuan Li^{1†}, Qi Yan¹, Lin Li¹, Yunfeng Hao¹, Dan Yao¹, Huaxing Si¹, Yingjun Zhao^{2*} and Wei Zhang^{1*}

Abstract

Background: Autoimmune encephalitis (AE) with neuronal surface antibodies (NSAbs) presents pathogenesis mediated by B cell-secreting antibodies. Rituximab is a second-line choice for the treatment for AE with NSAbs, which can cause B cell depletion via targeting CD20. However, the optimal protocol and dosage of rituximab combined with first-line therapy for NSAbs-associated AE remains unclear so far. In this study, we explored the efficacy and safety of low-dose rituximab combined with first-line treatment for NSAbs-associated AE.

Methods: Fifty-nine AE patients with NSAbs were enrolled, and retrospectively divided into common first-line therapy (41 patients) and combined low-dose rituximab (100 mg induction weekly with 3 circles, followed by 100 mg reinfusion every 6 months) with first-line therapy (18 patients). Outcome measures included changes in the Clinical Assessment Scale for Autoimmune Encephalitis (CASE) score (primary endpoint), changes in the modified Rankin Scale (mRS), the Mini-mental State Examination (MMSE), the patient and caregiver Neuropsychiatric Inventory (NPI) score at each visit (baseline, discharge, 6 months, 12 months and last follow-up) between two groups (secondary endpoint), as well as oral prednisone dosage, relapse and adverse effects during follow-up.

Results: Compared with traditional first-line therapy group, for primary outcome, CASE scores at last follow-up were significantly improved in combined rituximab group, as well as markedly improving changes of CASE scores between baseline and each visit. While changes of mRS, MMSE and NPI scores, as secondary endpoint, were all markedly accelerating improvement between baseline and each visit, as well as both oral prednisone dosage and relapse were also greatly reduced during follow-up. Meanwhile, longitudinal analysis in combination of rituximab cohort also revealed persistently marked amelioration in a series of scales from baseline even more than 1 year. Moreover, analysis in rituximab subgroup showed no difference in any clinical outcomes between combination with single first-line and with repeated first-line treatment (≥ 2 times), while compared to delayed combination with rituximab (> 3 months),

[†]Ying Du, Chao Zhao, Juntong Liu and Chuan Li contributed equally to this work

*Correspondence: yjzhao@xmu.edu.cn; tdzw@fmmu.edu.cn

¹ Department of Neurology, Tangdu Hospital, Fourth Military Medical University, Xi'an City 710038, Shaanxi Province, China

² Fujian Provincial Key Laboratory of Neurodegenerative Disease and Aging Research, Institute of Neuroscience, School of Medicine, Xiamen University, Xiamen 361005, China



© The Author(s) 2022, corrected publication 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

early initiation of combination (≤ 3 months) might achieve better improvements in CASE and MMSE assessment even 1 year later. No rituximab-correlated serious adverse events have been reported in our patients.

Conclusions: Our simplified regimen of combined low-dose rituximab firstly showed significantly accelerating short-term recovery and long-term improvement for AE with NSAbs, in parallel with markedly reduced prednisone dosage and clinical relapses. Moreover, opportunity of protocol showed earlier initiation (≤ 3 months) with better long-term improvement.

Keywords: Autoimmune encephalitis, Neuronal surface antibody, Rituximab, Low dose, Combined treatment, Clinical outcome

Introduction

Autoimmune encephalitis (AE) is a new spectrum of immune-mediated disorders in central nervous system (CNS), characterized by pathogenic autoantibodies against neuronal surface or intracellular proteins [1]. Up to now, autoantibodies against neuronal proteins, especial neuronal surface antibodies (NSAbs), have been linked to more than 15 AE subtypes, such as anti-N-methyl-D-aspartate receptor (NMDAR)-AE, anti-leucine-rich glioma-inactivated-1 (LGII)-AE and anti-contactin-associated protein-like-2 (CASPR2)-AE, making up the majority of seropositive AE subtypes [2]. Among them, most common NMDAR-AE, predominating in young women and children, mainly led to psychiatric symptoms, amnesia, epileptic seizures, reduced levels of consciousness and abnormal movements [3]. LGII-AE, particularly affecting middle-aged or elderly patients, frequently caused confusion, short-term memory deficits, faciobrachial dystonic seizures and hyponatremia [4]. CASPR2-AE is mostly observed in elderly men, usually presented LGII-like encephalitis and peripheral nerve hyperexcitability (neuromyotonia and myokymia) [5]. Despite heterogeneity of clinical features mediated by different antibodies in AE, the characterized manifestations were always involved in psychosis, cognitive impairments and seizures, far beyond motor dysfunctions [6]. NSAbs against surface antigens, such as NMDAR, LGII and CASPR2, may directly affect the targeted protein and cause clinical disturbances by blocking functions, interfering with synaptic protein interactions, or subsequent alterations of synaptic density. Furthermore, the underlying autoimmune processes may also lead to irreversible structural damages, as well as severe, progressive and refractory symptoms [7].

Treatment options for AE with NSAbs, ranging from broadly immunosuppressive agents to those targeting antibody-mediated pathogenesis, are mainly focused on achieving both better outcomes and fewer relapses. Common first-line immunotherapeutic agents for acute treatment include methylprednisolone pulse therapy (MPPT), intravenous immunoglobulin (IVIG), plasma exchange (PLEX) or combinations with less specificity

for pathogenesis, presenting short-term efficacy during administration. The addition of a second-line agent such as rituximab, cyclophosphamide or combinations will be initiated, if there is no meaningful clinical or radiological response to optimized first-line treatment after 2–4 weeks [8]. Then, a steroid-sparing therapy or a bridging strategy of oral prednisone with a gradual taper overlapping with azathioprine or mycophenolate mofetil (MMF) after completing acute therapy, should be implemented for sustained immunosuppression [9]. There are no established guidelines for AE with NSAbs treatment so far, and the traditional protocol is often complicated and empirically performed according to patient status and clinician opinion. Given the balance between efficacy and safety, an aggressive and practicable approach of immunotherapy after definite diagnosis is critical for AE patients [10].

Recently, a retrospective study based on real-world data has indicated that high-dose rituximab (at least 1 g once) is the most frequent second-line immunosuppressive agent used in AE with NSAbs, but fails to present significant privilege in combination with first-line treatment because of more severity at baseline and complicated addition of drugs in the rituximab cohort, according to scores of modified Rankin Scale (mRS) [11]. Moreover, mRS, initially designed to measure motor function for stroke, has limitations for assessing non-motor deficits of AE mainly presenting psychosis, amnesia and seizures, thereby probably misleading objective evaluation of clinical outcomes. Therefore, a new specialized scale, named the Clinical Assessment Scale for Autoimmune Encephalitis (CASE), has been developed to rate the severity of AE comprehensively [12], and validated for AE with NSAbs in Chinese patients, presenting more sensitive to clinical changes than mRS [13, 14]. Meanwhile, up to now, the infusion regimen, optimal dosage and clinical benefit of combined rituximab for AE with NSAbs treatment still need to be elucidated. Here, we performed a retrospective study to assess the clinical outcomes for patients with NSAbs-associated AE, who were treated with common first-line medications only or with first-line medications and low-dose rituximab (an induction with 100 mg

rituximab once a week for 3 cycles, followed by reinfusion 100 mg every 6 months at least for 1 year). We used improvement of CASE scores as the primary endpoint, and alterations in mRS, the Mini-mental State Examination (MMSE), the patient and caregiver Neuropsychiatric Inventory (NPI) scores as the secondary endpoints. We also evaluated clinical relapses, glucocorticoid reduction and adverse effects from different treatments. Without special notation, we will use the term “AE” to refer to AE with NSAbs only in the present study.

Materials and methods

Standard protocol approvals

This study was performed according to the Declaration of Helsinki, and approved by the Ethical Committee of Tangdu Hospital, Fourth Military Medical University. Moreover, we have provided patients and their relatives detailed information about the disease, and obtained the consent of the patients or their legal representatives to conventional first-line therapy and low-dose rituximab treatment, while written informed consent was obtained from all patients or their legal representatives.

Study population

The study recruited 72 Chinese patients with NSAbs-associated AE diagnosed in the Department of Neurology of Tangdu Hospital from April 2015 to April 2021. Finally, 59 patients were retrospectively collected and analyzed, while the other 13 were excluded due to incomplete data or lost follow-up. Among them, 18 patients with combined low-dose rituximab met the following inclusion criteria: (1) patients with detection of NMDAR-, LGI1- or CASPR2- antibodies in CSF and/or serum, and definite diagnosis of NSAbs-associated AE according to published criteria [15]; (2) mRS scores ≥ 3 [16] or CASE scores ≥ 5 [13] at baseline screening; (3) any documented treatment with low-dose (100 mg once) rituximab, and available information on the number and timing of infusions; (4) patients with combined low-dose rituximab received prior first-line immune treatments, including MPPT 1000 mg daily for 5 days and/or IVIG 2 g/kg over 5 days (0.4 g/kg/day). The exclusion criteria were: (1) combination with other antibodies against neuronal and glia antigens; (2) disease complicated by potentially acute or chronic viral or bacterial infections, such as HIV, latent hepatitis B, tuberculosis, syphilis, viral encephalitis and so on; (3) presence of other severe neurological or psychiatric complications, such as brain tumor, stroke, myasthenia gravis and so on. In addition, a control cohort of total 41 patients with only first-line immunotherapy were also enrolled, with the consistent inclusion and exclusion criteria.

Study design

All the patients had received at least one cycle of first-line immunotherapy defined as intravenous MPPT 1000 mg daily for 5 days, and/or IVIG 2 g/kg over 5 days (0.4 g/kg/day), followed by oral prednisone 30–60 mg/day for sustained immunosuppressive treatment with gradually tapering off [8, 9]. The combined regimen of rituximab was an induction of 100 mg once a week for 3 cycles, followed by reinfusions (100 mg once) at regular intervals (every 6 months) [17]. Relapse during follow-up was defined as new onset or worsening of encephalitis symptoms occurring after at least 2 months of improvement or stabilization [18], and judged by treating neurologists (Y Du, C Zhao and W Zhang) according to overall clinical impression. Simultaneously, detailed clinical status and lab examinations of each patient were evaluated at baseline and continuous 4 visits (discharge, 6 months, 12 months and last follow-up). Data on any immunotherapy and side effects of rituximab were recorded.

The primary efficacy endpoint of this study was the significant difference in the CASE score or accelerating improvement at each visit between combined low-dose rituximab treatment and common first-line therapy group. The secondary outcome measures were the marked differences in the mRS score, the MMSE score, the patient and caregiver NPI score or their accelerating improvements at each visit between two cohorts, as well as doses of oral prednisone and occurrence of relapses during the follow-up.

Laboratory detection of neuronal surface antibodies

Antibody testing was performed by cell-based assays (CBAs) and confirmed by immunofluorescence (commercial test kit panels Euroimmun, Lübeck) for NMDAR, LGI1, and CASPR2. Patients reached the following inclusion antibody criteria: NMDAR antibody was detected in serum by CBA ($>1:100$), followed by confirmation from immunofluorescence (in the absence of confirmatory immunofluorescence in serum, only CBA serum titers $>1:320$ were considered specific) and/or CSF positive; LGI1 antibody at any titer in CSF and/or serum; CASPR2 antibody $>1:100$ in serum and/or CSF positive [19]. Only IgG antibodies were considered relevant.

Clinical assessment of immunotherapy profiles

As described previously, the CASE scale, evaluated for the current status of AE, consists of nine items, including seizure (current time), memory dysfunction, psychiatric symptoms (delusion, hallucination, disinhibition, aggression), consciousness, language problem, dyskinesia/dystonia, gait instability and ataxia, brainstem dysfunction, and weakness. The total maximum score was 27.

Each item was based on a 3-point grading system, with the exception of the item “brainstem dysfunction,” which consisted of gaze paresis, tube feeding, and ventilator care due to hypoventilation [20]. Specifically, the items of “memory” and “language problem” were assessed mainly by communications and observations, rather than neurological examinations; the item of “seizure” was scored as 1 for controlled seizures with no need of dose-up, and scored as 2 for intractable seizures with the need of dose-up; in comatose patients, the items of “seizure,” “dyskinesia/dystonia,” and “brainstem dysfunction” could be used for evaluation, whereas all others were scored as 3 [13]. Moreover, two neurologists (J Liu and C Li) who were blinded to the diagnosis evaluated the scales independently by studying the detailed medical records, retrospectively. C Zhao repeated the assessment 1 month later.

The mRS scale consists of six grades (0–5 points), and predominantly captures the impact of motor deficits on functional independence [21]. The MMSE scale consists of 30 questions with the highest 30 points, and tests five cognitive domains, including time and place orientation (10 points), memory registration (3 points) and recall (3 points), attention and calculation (5 points), language and praxis (9 points), indicating higher scores with better cognition. Specifically, $MMSE \geq 27$ is considered normal, $26 \geq MMSE \geq 21$ is considered mild, $20 \geq MMSE \geq 10$ is considered moderate, while $MMSE \leq 9$ is considered severe cognitive impairment [22]. The NPI scale consists of 12 items of neuropsychiatric disturbances common in dementia, including delusions, hallucinations, agitation, dysphoria, anxiety, apathy, irritability, euphoria, disinhibition, aberrant motor behavior, night-time behavior disturbances, and appetite and eating abnormalities. The severity and frequency of each symptom are rated on the scripted questions for the patient’s caregiver, as well as assessment of caregiver distress by each neuropsychiatric disorder, then followed by a calculation of total NPI and total caregiver distress score [23]. All subjects were routinely assessed with mRS, MMSE, patient and caregiver NPI by two neurologists (D Yao and L Li) at baseline and each visit.

To adjust the possible differences in baseline functional status, the parameter “favorable clinical response” was analyzed at each visit under the following definition: improvement of the CASE scores by ≥ 5 points or achievement of the CASE scores ≤ 2 (as good) [20], improvement of the mRS scores by ≥ 2 points or achievement of the mRS scores ≤ 2 (as good) [21], or improvement of the MMSE scores by ≥ 10 points or achievement of the MMSE scores ≥ 27 (as good) [22].

Statistical analyses

Statistical tests were performed using GraphPad Prism 8.0 (GraphPad Software, Inc., San Diego, CA) and SPSS

version 26.0 (IBM, Armonk, NY, USA). Quantitative data with normal distributions were presented as mean \pm SD. Continuous variables conformance to skew distributions such as CASE, mRS, MMSE, NPI and the difference of scale score before and after rituximab were described as medians with the interquartile range (IQR) and analyzed with a Wilcoxon signed-rank test and a Wilcoxon rank sum test. Symptoms and demographic data were analyzed using the χ^2 test or Fisher exact test for categorical variables and Mann–Whitney *U* test for continuous variables. The time-weighted average prednisolone dose (mg/day) was calculated for each 4-week period after initial immunotherapy. The time-weighted average dose over each 4-week period was chosen to reflect inter and intraindividual variations in the interval and degree of dosage changes. One-way repeated measures analysis of variance (ANOVA) was conducted to analyze the effect of rituximab on the time-weighted average prednisolone dose. $p < 0.05$ was considered as significant.

Results

Patient characteristics with first-line or combined rituximab treatment

According to the designed inclusion and exclusion criteria, we identified 59 patients (26 females, 33 males) with NSAbs-associated AE, including 41 NMDAR-AE, 12 LGI1-AE, and 6 CASPR2-AE patients. Among them, our study cohort comprised 18 patients in the combined rituximab cohort (14 with NMDAR-AE, 3 with LGI1-AE, and 1 with CASPR2-AE), 41 patients with only first-line therapy in the control cohort (27 with NMDAR-AE, 9 with LGI1-AE, and 5 with CASPR2-AE), and no patient suffered from tumor during study. Gender, age at onset, duration from onset to diagnosis, follow-up duration, CSF/MRI/EEG profiles, and CSF/serum antibody profiles showed no difference between two groups. The clinical symptoms of AE presented heterogeneous, including seizures, cognitive impairments, psychiatric symptoms, decreased consciousness, autonomic dysfunction, movement disorder and fever, and also showed no difference between two groups at baseline (Table 1).

Compared to control group with only first-line therapy, patients with combined low-dose rituximab received less MPPT (72.2% vs 97.6%, $p = 0.008$) but more IVIG (94.4% vs 46.3%, $p < 0.0001$), while no difference was observed for combination of both first-line treatments between two groups (66.7% vs 43.9%, $p = 0.092$). The combined regimen of low-dose rituximab for 18 patients was an induction of 100 mg rituximab once a week for 3 cycles, followed by reinfusions (100 mg once) at regular intervals (every 6 months). In total combined rituximab cohort, median time from rituximab therapy was 74.5 days,

Table 1 Characterization of the patient cohort

	TOTAL			NMDAR-AE			LGI1-AE			CASPR2-AE		
	RTX (n = 18)	Ctrl (n = 41)	pvalue	RTX (n = 14)	Ctrl (n = 27)	pvalue	RTX (n = 3)	Ctrl (n = 9)	pvalue	RTX (n = 1)	Ctrl (n = 5)	pvalue
Gender:	8/10	18/23	.595	7/7	11/16	.406	1/2	5/4	.100	0/1	2/3	/
Female/Male												
Age at onset, y; mean (95% CI)	41.88(33-50)	37.61(32-42)	.356	35.28(27-42)	34.22(28-40)	.823	63.33(48-78)	52.88(43-62)	.190	70(/)	28.40(19-37)	/
From onset to diagnosis, d; median (IQR)	20(78)	24(52)	.863	14.5(23.75)	16(51)	.185	35(/)	60(63)	.864	190(/)	31(41)	/
Follow-up duration d; mean (95% CI)	1058(768-1349)	1362(1136-1587)	.120	1084(802-1376)	1488(1215-1556)	.059	1086(/)	1640(1278)	.926	244(/)	561(640)	/
Symptoms; n(%)												
Seizures	66.7%	61.0%	.455	71.4%	66.7%	.100	66.7%	55.6%	.100	0.00%	40%	/
Cognitive impairment	94.4%	85.4%	.422	92.9%	85.2%	.645	100%	100%	/	100%	60%	/
Psychiatric symptoms	83.3%	87.8%	.690	78.6%	88.9%	.393	100%	88.9%	.100	100%	80%	/
Decreased con- sciousness	44.4%	29.3%	.201	57.1%	33.3%	.189	0.0%	11.1%	.100	0.0%	40%	/
Autonomic dysfunction	27.8%	22.0%	.742	28.6%	25.9%	.100	33.3%	0.0%	.250	0.0%	40%	/
Movement disorder	44.4%	26.8%	.151	50%	22.2%	.089	33.3%	33.3%	.100	0.0%	40%	/
Fever	44.4%	39.0%	.457	57.1%	51.9%	.504	0.0%	11.1%	.100	0.0%	20%	/
CSF/MRI/EEG profiles												
CSF cc; median, (IQR)	7(67)	8(23)	.613	8.5(90.5)	12(26)	.264	4(/)	2(4)	.0282	2(/)	1(62)	
CSF protein; mean, (95% CI)	404.45(318-490)	353.82(288-419)	.370	372.05(274-469)	405.57(313-497)	.637	470.28(93-847)	247.21(179-314)	.009	660.7(/)	266.32(144-387)	/
CSF pressure; mean, (95% CI)	174.72(146-203)	161.46(140-182)	.463	182.85(147-218)	170.18(140-199)	.587	145.00(46-243)	131.66(100-162)	.632	150(/)	168.00(121-214)	/
MRI abnormali- ties; n (%)	50.0%	56.1%	.440	64.3%	51.9%	.336	0.0%	77.8%	.045	0.0%	40%	/
EEG abnormal- ities; n (%)	61.1%	43.9%	.175	71.4%	48.1%	.137	33.3%	33.3%	.100	0.0%	40%	/
CSF/Serum Ab profiles												
CSF Ab posi- tive	94.4%	80.5%	.252	100%	88.9%	.539	100%	77.8%	.100	0.0%	40%	/

Table 1 (continued)

TOTAL	NMDAR-AE			LGI1-AE			CASPR2-AE		
	RTX (n = 18)	Ctrl (n = 41)	pvalue	RTX (n = 14)	Ctrl (n = 27)	pvalue	RTX (n = 3)	Ctrl (n = 9)	pvalue
Serum Ab positive	38.9%	61.0%	.100	28.6%	48.1%	.321	66.7%	77.8%	.100
Both Ab positive	33.3%	41.5%	.386	28.6%	37.0%	.734	66.7%	55.6%	.100
Prior 1st-line immunotherapy; n (%)	100%	100%	/	100%	100%	/	100%	100%	/
1st-line therapy, n (%)									
MPPT	72.2%	97.6%	.008	85.7%	93.3%	.265	33.3%	100%	.045
IVIg	94.4%	46.3%	<.0001	92.9%	55.6%	.031	100%	33.3%	.182
Both	66.7%	43.9%	0.092	78.6%	51.9%	.176	33.3%	33.3%	.100
RTX therapy									
Time from RTX therapy, d; median (IQR)	74.5(410)	/	/	128.5(405)	/	/	6(/)	/	/
No. of RTX infusions, n; median (IQR)	5(4)	/	/	5(3)	/	/	3(/)	/	/
Cumulative RTX dose, g; median (IQR)	500(400)	/	/	500(300)	/	/	300(/)	/	/
1st to last infusion, d; median (IQR)	389(864)	/	/	457.5(863)	/	/	15(/)	/	/
Averaged dose of Prednisone, mg/day; median (IQR)	4.375(7.67)	27.13(21.67)	<.0001	5.563(4.011)	28.89(21.67)	<.0001	0	10.83(19.52)	.009
Relapses; n (%)									
After 1st-line therapy	33.3%	12.2%	.074	42.9%	18.5%	0.140	33.3%	0.0%	.250
After RTX therapy	0	/	/	0	/	/	0	/	/

AE autoimmune encephalitis, NMDAR N-methyl-D-aspartate receptor, LGI1 leucine-rich glioma-inactivated-1, CASPR2 contactin-associated protein-like-2, RTX rituximab, Ctrl control, CI confidence interval, IQR interquartile range, CSF cerebrospinal fluid, MRI magnetic resonance imaging, EEG electroencephalogram, cc cell count, Ab antibody, MPTP methylprednisolone pulse therapy, IVIg intravenous immunoglobulin. *p* values reaching statistical significance are indicated in bold

median number of rituximab infusions was 5 times, median dose of cumulative rituximab was 500 mg, and median time from initiation to last infusion was 389 days (Table 1).

Clinical outcomes

Compared to control cohort without rituximab, patients in combined rituximab cohort presented a more severe tendency in CASE, mRS, MMSE, patient or caregiver NPI score at baseline, but without significantly statistical differences ($p > 0.05$). However, at last follow-up, combined rituximab patients still showed a better outcome of CASE score than that of control group ($p = 0.037$). Moreover, in order to adjust the possible influence of baseline status, we also evaluated the improvement from baseline of CASE score at each visit, and the results still showed that patients with rituximab presented a significant improvement from baseline of CASE scores than those without rituximab at discharge ($p = 0.01$), 6 months ($p = 0.016$), 12 months ($p = 0.013$) and last follow-up ($p = 0.001$), respectively, suggesting the achievement of primary endpoint as designed in our study. Meanwhile, in the secondary outcome measures, clinical response reflected by a series of scales, including mRS, MMSE, patient and caregiver NPI, was noted at each visit. After adjusting the possible influence by baseline status, compared with control group, mRS, MMSE, patient and caregiver NPI also showed significant improvement from baseline in those with combined rituximab from discharge, and almost lasting till the study end, respectively (Table 2) (Fig. 1).

Furthermore, in the analysis of parameter for “favorable clinical response”, compared with control group, combined rituximab group presented better percentage in improvement of CASE scores at discharge (77.8% vs 41.5%, $p = 0.01$) and 12 months (94.4% vs 63.4%, $p = 0.011$) (Fig. 2A), as well as higher frequency in improvement of mRS scores at discharge (72.2% vs 41.5%, $p = 0.028$) (Fig. 2B) and MMSE scores at last follow-up (94.4% vs 70.7%, $p = 0.039$) (Fig. 2C). Meanwhile, compare to control cohort, the cumulative oral prednisone doses obviously decreased in rituximab cohort ($p < 0.001$) (Fig. 2D), as well as the time-weighted average prednisolone dosage in rituximab cohort was also markedly reduced than that in control cohort within the follow-up ($p < 0.001$) (Fig. 2E). Especially, during the study, 12 patients in total 59 patients had 12 relapses (11 with NMDAR-AE, 1 with LGI1-AE). Among them, 7 relapses occurred before combined rituximab initiation in rituximab cohort (6 with NMDAR-AE, 1 with LGI1-AE), and no relapses occurred after rituximab treatment, while the other 5 relapses occurred in the control cohort (5 with

NMDAR-AE). Cumulatively, there were 12 relapses after only first-line treatment in total AE patients, which was showed significantly frequent than that observed after initiation of rituximab (20.33% vs 0%, $p = 0.037$) (Fig. 2F).

Treatment response and follow-up in each cohort

Further analyses about each cohort, including control group with only first-line therapy and combined rituximab group, were also performed for longitudinal changes in CASE, mRS, MMSE, patient and caregiver NPI scores, respectively. In both control cohort and combined rituximab cohort, scores of CASE, mRS, MMSE, patient and caregiver NPI were all greatly improved from baseline at each visit (Additional file 1: Table S1). When we analyzed the CASE (≤ 2), mRS (≤ 2) or MMSE (≥ 27) scores for favorable outcomes in the rituximab cohort throughout follow-up in more detail, we found that patients with rituximab showed significant improvement in CASE scores at discharge ($p = 0.001$) and 12 months ($p = 0.049$) (Fig. 3A), continuously improving trend in mRS scores with achievement of statistical significance at discharge ($p < 0.0001$) and 6 months ($p = 0.015$) (Fig. 3B), and sustained improving trend in MMSE scores with achievement of statistical significance at discharge ($p = 0.015$) and last visit (> 12 months) ($p = 0.004$) (Fig. 3C). While in analysis of a series of scales in the control cohort without rituximab during follow-up, we also found that patients in control group presented significant improvement in CASE scores at discharge ($p < 0.0001$) and 12 months ($p = 0.039$), great amelioration in mRS scores at discharge ($p < 0.0001$) and 6 months ($p < 0.0001$), and marked promotion in MMSE scores only at 6 months ($p < 0.0001$). Altogether, these results proposed that both only first-line and combined rituximab therapy showed significantly comprehensive efficiency within 1 year by a series of scales assessment, while combined rituximab treatment seemed to especially present sustained improvement in cognitive impairment even 1 year later.

Clinical analysis for subgroup in rituximab cohort

In our study, all the 18 patients with rituximab received at least one cycle of prior first-line immunotherapy, defined as intravenous methylprednisolone 1000 mg daily for 5 days, and/or IVIG 2 g/kg over 5 days (0.4 g/kg/day). Among them, 9 patients were administrated first-line treatment only for one cycle, while the other 9 patients received repeated first-line therapy more than twice. Moreover, 10 patients in rituximab cohort were initiated rituximab within 3 months, while the other 8 patients were delayed for initiation over 3 months. Further analysis was performed to assess the possible influence of repeated first-line treatment and rituximab initiating

opportunity on clinical outcome in different subgroups of rituximab cohort (Additional file 2: Table S2 and Additional file 3: Table S3). The results showed clearly that compared to single first-line treatment combined with low-dose rituximab subgroup, repeated first-line administration combined with rituximab presented no significant difference in CASE (Fig. 4A), mRS (Fig. 4B), MMSE (Fig. 4C), patient or caregiver NPI evaluation at baseline and each visit (Additional file 2: Table S2). However, compared to the subgroup with delayed initiation of rituximab over 3 months, early initiation of rituximab within 3 months led to a significant improvement in CASE ($p=0.039$) (Fig. 4D) and MMSE ($p=0.046$) (Fig. 4F) assessment at last follow-up, respectively, but still showed no significance in mRS scores (Fig. 4E), patient or caregiver NPI at baseline and each visit (Additional file 3: Table S3). Altogether, these results suggested that repeated first-line treatment presented no influence on clinical outcomes in patients with combination of rituximab, while early initiation of combined rituximab within 3 months seemed to showed significant advantages in long-term clinical improvement.

Adverse effects and safety of rituximab

In a total of 18 patients on rituximab, two had infusion-related symptom which presented as skin rash, fever during the administration of rituximab. However, the symptoms gradually disappeared after oral cetirizine. Severe infusion adverse events did not occur in all patients. Of note, the side effects of glucocorticoids as central obesity, dropsy, acne, osteoporosis, abnormal glucose tolerance, hypertension, psychiatric disorder were often presented in non-rituximab cohort for long-term large oral doses of hormones.

Discussion

In present study, the 3 main findings about rituximab treatment for AE are that: (1) our simplified regimen of low-dose rituximab (100 mg once) combined with common first-line therapy significantly accelerates comprehensive short-term recovery within 1 year, as well as markedly contributing to long-term improvement even after at least 1 year. (2) Our refined protocol of rituximab infusions leads to faster oral prednisolone gradual taper and withdrawal, in parallel with markedly sustained clinical remission and reduced relapses. (3) Opportunity of rituximab schedule shows earlier initiation with better improvement, while frequency of first-line therapy has no influence on satisfactory outcome with rituximab combination.

As we know, self-reactive B cells are subject to the processes of negative selection for elimination, such as deletion, receptor editing and induction of anergy,

throughout the development in the bone marrow and spleen [24]. However, emerging data suggest defective B cell tolerance checkpoints in several AE (such as NMDAR-AE, LGI1-AE and CASPR2-AE), increasing autoreactive immature B cells that are not removed but can be activated and enter germinal centers [25, 26]. Several mechanisms may contribute to the loss of B cell tolerance in peripheral lymph nodes in the context of tumor ectopic expressions or potential viral infections, particularly by inducing B cell-intrinsic Toll-like receptor (TLR) signal together with B cell receptor (BCR) ligation, or activating T helper (Th) cell with same antigen stimulation, thereby leading to consecutive B cell clonal expansion, class switch, affinity maturation and NSAbs production [24, 27, 28]. Subsequently, activation of innate immune-mediated cytokines and TLR ligands leads to disruption of the blood brain barrier (BBB), allowing infiltration of autoreactive memory B and plasma cell, followed by proliferation with B cell activating factor (BAFF), and releasing large amounts of NSAbs in CNS [29]. Among them, anti-NMDAR antibodies are mainly IgG1 subclass, leading to rapid reduction of neuronal surface NMDAR by cross-linking, internalization and degradation, while antibodies against LGI1 and Caspr2 are predominantly IgG4, inducing neuronal dysfunction by interrupting the trans-synaptic binding of LGI1 or Caspr2 to its postsynaptic receptor a disintegrin and metalloprotease 22 (ADAM22) and likewise ADAM23 at the presynaptic site, thus causing a series of clinical phenotypes [30, 31].

Rituximab exerts therapeutic effect through its target, cluster of differentiation 20 (CD20), an integral membrane protein mainly expressed on B lymphocytes. During the autoimmune pathological process, the expression of CD20 is progressively increased in B cells at different developmental stages and sustainedly expressed at a high level on the surface of antibody-specific memory B cells and plasmablasts [32]. In NSAbs-associated AE, CD20 may act as a physically coupled link to BCR and other surface molecules or cytoplasmic proteins, such as major histocompatibility complex class II (MHCII), CD40 molecules, and tyrosine kinases, thereby regulating cell cycle progression and proliferation of B lymphocytes [33]. Moreover, activated CD20+B cells can present same specific antigens to T lymphocytes in association with MHC molecules in the presence of various costimulatory factors, thus promoting T cells activation and differentiation. Subsequently, these T cells can produce a variety of cytokines and chemokines to regulate the maturation and migration of peripheral immune effectors, such as Th, CD8+ T and myeloid cells, secreting a range of pro-inflammatory mediators to induce neuroinflammation within the CNS parenchyma [34]. Therefore, rituximab

Table 2 Comparison of clinical outcomes between rituximab cohort and control cohort

Evaluation scales	Baseline			1st visit			2nd visit			3rd visit			Last visit		
	RTX	Ctrl	p	RTX	Ctrl	p	RTX	Ctrl	p	RTX	Ctrl	p	RTX	Ctrl	p
CASE	Scores median(IQR)	8 (7.25)	6 (2)	0.079	3 (4.5)	3 (2)	0.763	2 (3.25)	2 (2)	0.834	1.5 (1.25)	2 (2)	0.172	0 (1.25)	1 (2)
	Scores differ ± median (IQR)	-	-	-	-5 (4.5)	-3 (2)	0.010	-6.5 (5.5)	-3 (3)	0.016	-6.5 (7.3)	-4 (3)	0.013	-7 (7)	-4 (3)
mRS	Scores median(IQR)	4 (1.25)	4 (1)	0.083	2 (1)	3 (1)	0.270	2 (1)	2 (1.5)	0.771	1 (1.25)	1 (1)	0.532	0.5 (1.25)	1 (2)
	Scores differ ± median (IQR)	-	-	-	-1 (1)	-1 (0.5)	0.010	-2 (1.0)	-2 (2)	0.140	-3 (2)	-2 (1)	0.043	-3.5 (2)	-3 (1)
MMSE	Scores median (IQR)	14.5 (16.25)	20 (15.5)	0.062	23.5 (11)	23 (10)	0.785	25.5 (7.75)	24 (8)	0.907	26 (6.25)	27 (7)	0.522	28.5 (3)	27 (5)
	Scores differ ± median (IQR)	-	-	-	+8 (8.8)	+3 (7.5)	0.006	+11 (15.5)	+4 (1.3)	0.046	+10.5 (16)	+5 (14)	0.037	+13 (15.3)	+7 (14)
Patient NPI	Scores median (IQR)	15 (20.5)	9 (10)	0.066	1.5 (4.5)	3 (5.5)	0.532	0 (4)	0 (4)	0.863	0 (2)	0 (3)	0.538	0 (0)	0 (1.5)
	Scores differ ± median (IQR)	-	-	-	-14 (14.3)	-6 (10)	0.021	-13 (16)	-7 (11.5)	0.044	-14 (20)	-8 (11.5)	0.038	-15 (20.8)	-8 (11.5)
Caregiver NPI	Scores median (IQR)	6.5 (5.75)	5 (5)	0.054	1 (3)	2 (3)	0.461	0 (2)	0 (2)	0.785	0 (0.25)	0 (1.5)	0.381	0 (0.25)	0 (1.5)
	Scores differ ± median (IQR)	-	-	-	-6 (5)	-2 (4)	0.006	-5 (7.3)	-3 (5.5)	0.043	-6.5 (7.8)	-4 (5)	0.038	-6.5 (6.3)	-3 (5)

Detailed clinical status was evaluated by a series of AE-associated scales at baseline before treatment and continuous 4 visits after treatment. 1st visit: at discharge, 2nd visit: 6 months later, 3rd visit: 12 months later, 4th visit: last follow-up with at least > 12 months

RTX rituximab cohort, Ctrl control cohort, CASE the Clinical Assessment Scale for Autoimmune Encephalitis, mRS the modified Rankin Scale score, MMSE the Mini-mental State Examination score, NPI the Neuropsychiatric Inventory, IQR interquartile rang, p values reaching statistical significance are indicated in bold





Fig. 1 Flowchart of study design. NSAbs: neuronal surface antibodies; AE: autoimmune encephalitis, NMDAR: anti-N-methyl-D-aspartate receptor; LGI1: anti-leucine-rich glioma-inactivated 1; CASPR2: anti-contactin-associated protein-like-2; CASE: the Clinical Assessment Scale for Autoimmune Encephalitis, mRS: the modified Rankin scale, MMSE: the Mini-mental State Examination, NPI: the Neuropsychiatric Inventory

may inhibit neuroinflammation via targeting CD20+B lymphocytes, which results in beneficial effects for AE treatment.

Although instances of spontaneous recovery without immunotherapy have also been reported occasionally in AE, the disease is mostly presented as a progressively monophasic process with rare recurrence (approximately 10–20%) but apparent sequelae, suggesting the irreversible neuronal damages, and advocating the necessity for prompt and persistent interruption of pathogenic immune activity [35]. Traditional AE first-line therapy, including corticosteroids, IVIg, and PLEX, has limitation of less specificity for pathogenesis or shorter maintenance for treatment, while oral prednisone for bridging


and steroid-sparing, azathioprine or MMF for sustained immunosuppression, both have common deficiency of less specificity, as well as complicated diverse regimen and continuous adverse effects [8, 9]. Proposals derived from recent systematic review for AE treatment are favored in early initiation of immunotherapy and addition of second-line agents, thereby resulting in better functional outcomes and lower relapses with manageable side effects [36]. Given the largely B cell-secreted antibody mediating the disease pathogenesis, it should be considered that a combination of immunotherapeutic agents targeting B cells may be urgently required for a more efficient regimen of AE treatment [37]. Therefore, rituximab is preferentially selected in second-line agents



due to substantially special efficacy and relatively reliable safety.

Rituximab as a second-line agent for AE, initially approved for treatment of lymphoma, is a human/murine chimeric monoclonal antibody directed against a differentiation glycoprotein CD20 participating in B lymphocytes activation and proliferation. After binding to CD20 on the B cells surface, rituximab specifically depletes target B lymphocytes (such as naïve B cells, memory B cells and some plasmablasts) by antibody-mediated cellular toxicity, complement activation and induction of apoptosis, thereby reducing B cells response, and causing therapeutic immunosuppression [38]. Currently, activation of self-reactive B cells and their subsequent proliferation and differentiation into auto-antigen reactive memory B cells and autoantibody-secreting plasma cells, play pivotal pathogenic roles in antibody-mediated neurological diseases, such as AE, neuromyelitis optica spectrum disorders (NMOSD) and myasthenia gravis (MG).


Therefore, the off-label use of rituximab for deleting the antigen-specific memory B-cell populations and hence preventing the formation of new plasmablasts which secrete the pathogenic antibodies, was gradually emphasized and presented potential advantages in AE treatment [34]. Consensus criteria on the appropriate time to initiate a second-line agent such as rituximab are yet to be established in AE, but a quick procession is favored, regardless of the response to first-line therapy. When rituximab is used since the acute setting, it may have the added benefit of a potentially faster onset of action, and also serve as a bridging therapy to prevent early relapses that might happen if immunosuppression is abruptly discontinued, as well as an optimal alternative for a sustained immunosuppressant [18]. Thus, the previously complex regimen for AE might be simplified and refined as combination of regular rituximab infusions with conventional first-line therapy.



In reference to a variety of researches about rituximab treatment in AE, there is great heterogeneity concerning dosages. Currently, the empirical protocols for AE are mainly derived from other disease processes such as lymphoma or rheumatoid arthritis (RA), including 375 mg/m² weekly for 4 consecutive weeks or two doses of 1000 mg 2 weeks apart, then followed by reinfusions at fixed intervals for immunosuppressive maintenance, because of circulating B cell below the detectable range for 6–8 months after administration [39]. Moreover, the optimal dosage of rituximab for balancing between safety and efficacy are still ambiguous in AE treatment, and empirical off-label attempt primarily comes from high-dose therapy for lymphomas, usually exerting more medical expenses and serious adverse events [40]. Indeed, the dysfunctional B cells commonly present with normal circulating count in autoimmune diseases, which is different from the high tumor burden in lymphomas, and low-dose rituximab seems to be sufficient and effective for complete depletion of peripheral CD20⁺B cells [41]. Recently, reduced low-dose of 100 mg rituximab per infusion for treatment has been tried in some neurological autoimmune disorders, such as NMOSD, MG, multiple sclerosis (MS) and neuro-Behçet's disease (NBD), with the protocol of induction per week for 3

cycles, then followed by reinfusions at regular intervals. The approach still presented good responsive in depleting B cells, improving clinical symptoms and preventing relapses with favorable side-effect and medical cost [16, 42–44]. Thus, in present study, we performed a simplified regimen in AE treatment, including regular induction of 3 cycles for acute and bridging management, and subsequent reinfusions (100 mg once) at fixed interval (every 6 months) for sustained immunosuppression.

Meanwhile, the scales for evaluating clinical severity and therapeutic response in AE were also continuously improving and updating. Currently, because of no special tools for AE assessment, the modified Rankin scale (mRS) was widely applied to measure neurological severity and outcome [18, 20]. As we know, the mRS was primarily developed and mainly weighted for estimating prognosis of motor function in acute stroke management, while patients with AE usually presented a variety of symptoms beyond motor deficits, including behavioral changes, memory impairments, seizures, speech disorders, abnormal movements, decreased consciousness, and cerebellar ataxia, which might also interfere with each other in evaluation [12, 45]. Some other scales for special functional domains, such as MMSE for cognitive damages and NPI for neuropsychiatric symptoms, still



had obvious limitations of lack diversity in assessment for AE [22, 23]. Recently, prognostication and estimation tools specifically developed for AE, such as CASE scale with description in detail and validation in practice, might help to select those patients required for more aggressive immunotherapy, and comprehensively and accurately evaluate their clinical outcomes [12]. Hence, according to severity designed previously as mRS scores ≥ 3 or CASE scores ≥ 5 , we retrospectively chose moderate and severe AE patients received first-line therapy with or without combination of low-dose rituximab in this study, and also discussed changes of CASE, mRS, MMSE, patient and caregiver NPI scale as well as glucocorticoid dosage and relapses between different cohorts.

As we found during the follow-up, compared to control cohort with common first-line therapy for AE, combined low-dose rituximab cohort not only showed much better outcome in CASE scale evaluation even 1 year later, but also significantly accelerated improvement in CASE, mRS, MMSE, patient and caregiver NPI from baseline within 1 year, as well as markedly reducing occurrence of relapse and oral prednisone dosage, indicating the potential privilege of our simplified regimen of low-dose rituximab in both long-term and short-term prognosis, along

with sustained immunosuppression. Meanwhile, longitudinal self-control analysis in both groups also revealed continuously marked amelioration in a series of scales from baseline during at least 1 year, whereas the persistent improvement might be presented even more than 1 year in combination of rituximab. Moreover, further analysis in rituximab cohort showed no difference in any clinical outcomes between combination with single first-line and with repeated first-line treatment (≥ 2 times), while compared with delayed combination with rituximab subgroup (> 3 months), early initiation of combination (≤ 3 months) might achieve better improvements in CASE and MMSE assessments.

This study was limited by its uncontrolled design without comparison with natural course of AE or other dosing regimens, as well as retrospective observational analysis, small sample size and limited follow-up time, and a bias in selecting patients could not be completely ruled out. The data were collected during routine clinical practice rather than a formal study setting, which meant limitations of quality and quantity varied among patients. Meanwhile, we have altered AE treatment strategy when rituximab became a preferred choice for immunosuppression, causing only a small number of patients with

low-dose rituximab combination enrolled for therapeutic protocol. Since randomized trials are difficult to conduct in rare diseases such as AE, real-world data might contribute to important information on treatment profiles and protocols. Although limited data on simplified regimen of combined optimal low-dose rituximab in our study, the result might be encouraging and presenting therapeutic implications for AE.

Altogether, in present study, the simplified regimen of combined low-dose rituximab (100 mg once) with common first-line therapy for AE with NSAbs, to our knowledge, firstly showed effective for short-term and long-term improvement, in parallel with reduced immunosuppressant and relapses, suggesting the advantages and benefits for combination of low-dose rituximab in the disease course. Moreover, the opportunity of rituximab protocol showed earlier initiation with better improvement, while frequency of first-line treatment had no influence on satisfactory outcome with rituximab combination. Our reports may expand therapeutic options and provide helpful references for NSAbs-associated AE, and further studies to corroborate these findings are warranted.

Abbreviations

AE: Autoimmune encephalitis; CNS: Central nervous system; NSAbs: Neuronal surface antibodies; NMDAR: N-Methyl-D-aspartate receptor; LGI1: Leucine-rich glioma-inactivated-1; CASPR2: Contactin-associated protein-like-2; MPPT: Methylprednisolone pulse therapy; IVIG: Intravenous immunoglobulin; PLEX: Plasma exchange; MMF: Mycophenolate mofetil; mRS: Modified Rankin Scale; CASE: Clinical Assessment Scale for Autoimmune Encephalitis; NEOS: Anti-NMDAR Encephalitis One-Year Functional Status; MMSE: Mini-mental State Examination; NPI: Neuropsychiatric Inventory; CBAs: Cell-based assays; CSF: Cerebrospinal fluid; MRI: Magnetic resonance imaging; EEG: Electroencephalogram; TLR: Toll-like receptor; BCR: B cell receptor; BAFF: B cell activating factor; ADAM: A disintegrin and metalloprotease; CD20: Cluster of differentiation 20; MHCII: Major histocompatibility complex class II; RA: Rheumatoid arthritis; NMOSD: Neuromyelitis optica spectrum disorders; MS: Multiple sclerosis; NBD: Neuro-Behçet's disease; MG: Myasthenia gravis.

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12974-022-02622-8>.

Additional file 1: Table S1. Clinical improvements in rituximab cohort and control cohort

Additional file 2: Table S2. Comparison of clinical outcomes between single first-line and repeated first-line treatment subgroups in rituximab cohort

Additional file 3: Table S3. Comparison of clinical outcomes between early and delayed initiation of rituximab subgroups in rituximab cohort

Acknowledgements

We thank the patients and relatives for their collaboration.

Author contributions

YD, CZ, JL and CL contributed equally to this study and are co-first authors. WZ and YZ had full access to all the data in the study and takes full responsibility for the integrity of the data and the accuracy of the data analysis. Study concept and design: YD, CZ, WZ. Acquisition, analysis, or interpretation of data:

All authors. Drafting of the manuscript: YD, CZ, YZ, WZ. Critical revision of the manuscript for important intellectual content: all authors. Statistical analysis: YD, CZ, JL, CL. Study supervision: YZ, WZ. All authors reviewed the manuscript. All authors read and approved the final manuscript.

Funding

This work was supported by research Grants from the National Natural Science Foundation of China (81971003, 82171406, 82271457), Key project of research and development plan of Shaanxi Province (2018SF-086), Key project of basic natural science research program of Shaanxi Province (2019JZ-18), Tangdu hospital innovation development foundation (2018QYTS010, 2019QYTS002), Tangdu hospital clinical research project (2021LCYJ040), Tangdu hospital foundation for social recruitment talent (2021SHRC011).

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author.

Declarations

Ethics approval and consent to participate

We provided patients detailed information about the disease, and obtained the consent of the patients to repeated low-dose rituximab treatment. We have also reported to the Ethical Committee of Tangdu Hospital, Fourth Military Medical University, and obtained the approval from the committee.

Consent for publication

We obtained written informed consent from both patients for publication of this report and any accompanying images. A copy of the written consent is available for review by the Editor-in-Chief of this journal.

Competing interests

The authors declare that they have no competing interest.

Received: 5 July 2022 Accepted: 13 October 2022

Published: 22 October 2022

References

- Dalmau J, Graus F. Antibody-mediated encephalitis. *N Engl J Med*. 2018;378(9):840–51.
- Dalmau J, Geis C, Graus F. Autoantibodies to synaptic receptors and neuronal cell surface proteins in autoimmune diseases of the central nervous system. *Physiol Rev*. 2017;97(2):839–87.
- Dalmau J, Lancaster E, Martinez-Hernandez E, et al. Clinical experience and laboratory investigations in patients with anti-NMDAR encephalitis. *Lancet Neurol*. 2011;10(1):63–74.
- Irani SR, Alexander S, Waters P, et al. Antibodies to Kv1 potassium channel-complex proteins leucine-rich, glioma inactivated 1 protein and contactin-associated protein-2 in limbic encephalitis, Morvan's syndrome and acquired neuromyotonia. *Brain*. 2010;133(9):2734–48.
- Irani SR, Pettingill P, Kleopa KA, et al. Morvan syndrome: clinical and serological observations in 29 cases. *Ann Neurol*. 2012;72(2):241–55.
- Mader S, Brimberg L, Diamond B. The role of brain-reactive autoantibodies in brain pathology and cognitive impairment. *Front Immunol*. 2017;8:1101. <https://doi.org/10.3389/fimmu.2017.01101>.
- Prüss H. Autoantibodies in neurological disease. *Nat Rev Immunol*. 2021;21(12):798–813.
- Abboud H, Probasco JC, Irani S, Ances B, Benavides DR, Bradshaw M, Christo PP, et al. Autoimmune encephalitis: proposed best practice recommendations for diagnosis and acute management. *J Neurol Neurosurg Psychiatry*. 2021;92(7):757–68.
- Abboud H, Probasco J, Irani SR, Ances B, Benavides DR, Bradshaw M, et al. Autoimmune encephalitis: proposed recommendations for symptomatic and long-term management. *J Neurol Neurosurg Psychiatry*. 2021;92(8):897–907.
- Shin YW, Lee ST, Park KI, Jung KH, Jung KY, Lee SK, Chu K. Treatment strategies for autoimmune encephalitis. *Ther Adv Neurol Disord*. 2017;11:1756285617722347. <https://doi.org/10.1177/1756285617722347>.

11. Thaler FS, Zimmermann L, Kammermeier S, Strippel C, Ringelstein M, Kraft A, et al. Rituximab treatment and long-term outcome of patients with autoimmune encephalitis: real-world evidence from the GENERATE registry. *Neurol Neuroimmunol Neuroinflamm*. 2021;8(6): e1088. <https://doi.org/10.1212/NXI.0000000000001088>.
12. Lim JA, Lee ST, Moon J, Jun JS, Kim TJ, Shin YW, et al. Development of the clinical assessment scale in autoimmune encephalitis. *Ann Neurol*. 2019;85(3):352–8.
13. Cai MT, Lai QL, Zheng Y, Fang GL, Qiao S, Shen CH, Zhang YX, Ding MP. Validation of the clinical assessment scale for autoimmune encephalitis: a multicenter study. *Neurol Ther*. 2021;10(2):985–1000.
14. Zhang Y, Tu E, Yao C, Liu J, Lei Q, Lu W. Validation of the clinical assessment scale in autoimmune encephalitis in Chinese patients. *Front Immunol*. 2021;12: 796965. <https://doi.org/10.3389/fimmu.2021.796965>.
15. Graus F, Titulaer MJ, Balu R, et al. A clinical approach to diagnosis of autoimmune encephalitis. *Lancet Neurol*. 2016;15(4):391–404.
16. Qiao S, Wu HK, Liu LL, Zhang RR, Wang ML, Han T, Zhang SC, Liu XW. Characteristics and prognosis of autoimmune encephalitis in the east of China: a multi-center study. *Front Neurol*. 2021;12: 642078. <https://doi.org/10.3389/fneur.2021.642078>.
17. Yang CS, Yang L, Li T, Zhang DQ, Jin WN, Li MS, Su N, Zhangning N, Liu Q, Shao ZH, Yu C, Shi FD. Responsiveness to reduced dosage of rituximab in Chinese patients with neuromyelitis optica. *Neurology*. 2013;81(8):710–3.
18. Titulaer MJ, McCracken L, Gabioudo I, Armangué T, Glaser C, Iizuka T, et al. Treatment and prognostic factors for long-term outcome in patients with anti-NMDA receptor encephalitis: an observational cohort study. *Lancet Neurol*. 2013;12(2):157–65.
19. Bien CG, Bien CI, Dogan Onugoren M, De Simoni D, Eigler V, Haensch CA, et al. Routine diagnostics for neural antibodies, clinical correlates, treatment and functional outcome. *J Neurol*. 2020;267(7):2101–2014.
20. Lee WJ, Lee ST, Moon J, Sunwoo JS, Byun JI, Lim JA, et al. Tocilizumab in autoimmune encephalitis refractory to rituximab: an institutional cohort study. *Neurotherapeutics*. 2016;13(4):824–32.
21. Lee WJ, Lee ST, Shin YW, Lee HS, Shin HR, Kim DY, et al. Teratoma removal, steroid, IVIG, rituximab and tocilizumab (T-SIRT) in anti-NMDAR encephalitis. *Neurotherapeutics*. 2021;18(1):474–87.
22. Hang HL, Zhang JH, Chen DW, Lu J, Shi JP. Clinical characteristics of cognitive impairment and 1-year outcome in patients with anti-LGI1 antibody encephalitis. *Front Neurol*. 2020;11:852. <https://doi.org/10.3389/fneur.2020.00852>.
23. Chen W, Wang M, Gao L, Huang Z, Lin Y, Xue Q, Liu G, Zhang Y, Su Y. Neurofunctional outcomes in patients with anti-leucine-rich glioma inactivated 1 encephalitis. *Acta Neurol Scand*. 2021;144(6):632–9.
24. Rawlings DJ, Metzler G, Wray-Dutra M, Jackson SW. Altered B cell signaling in autoimmunity. *Nat Rev Immunol*. 2017;17(7):421–36.
25. Cyster JG, Allen CDC. B Cell Responses: Cell Interaction Dynamics and Decisions. *Cell*. 2019;177(3):524–40.
26. Nemazee D. Mechanisms of central tolerance for B cells. *Nat Rev Immunol*. 2017;17(5):281–94.
27. Kuraoka M, Snowden PB, Nojima T, Verkoczy L, Haynes BF, Kitamura D, Kelsoe G. BCR and endosomal TLR signals synergize to increase AID expression and establish central B cell tolerance. *Cell Rep*. 2017;18(7):1627–35.
28. Sanderson NS, Zimmermann M, Eilinger L, Gubser C, Schaeren-Wiemers N, Lindberg RL, Dougan SK, Ploegh HL, Kappos L, Derfuss T. Cocapture of cognate and bystander antigens can activate autoreactive B cells. *Proc Natl Acad Sci USA*. 2017;114(4):734–9.
29. Ding H, Jian Z, Stary CM, Yi W, Xiong X. Molecular pathogenesis of anti-NMDAR encephalitis. *Biomed Res Int*. 2015;2015: 643409. <https://doi.org/10.1155/2015/643409>.
30. Moscato EH, Peng X, Jain A, Parsons TD, Dalmau J, Balice-Gordon RJ. Acute mechanisms underlying antibody effects in anti-N-methyl-D-aspartate receptor encephalitis. *Ann Neurol*. 2014;76(1):108–19.
31. Kornau HC, Kreye J, Stumpf A, Fukata Y, Parthier D, Sammons RP, Imbrosci B, Kurpijuweit S, Kowski AB, Fukata M, Prüss H, Schmitz D. Human cerebrospinal fluid monoclonal LGI1 autoantibodies increase neuronal excitability. *Ann Neurol*. 2020;87(3):405–18.
32. Chen TX, Fan YT, Peng BW. Distinct mechanisms underlying therapeutic potentials of CD20 in neurological and neuromuscular disease. *Pharmacol Ther*. 2022;238: 108180. <https://doi.org/10.1016/j.pharmthera.2022.108180>.
33. Polyak MJ, Li H, Shariat N, Deans JP. CD20 homo-oligomers physically associate with the B cell antigen receptor. Dissociation upon receptor engagement and recruitment of phosphoproteins and calmodulin-binding proteins. *J Biol Chem*. 2008;283(27):18545–52.
34. Lee DSW, Rojas OL, Gommerman JL. B cell depletion therapies in autoimmune disease: advances and mechanistic insights. *Nat Rev Drug Discov*. 2021;20(3):179–99.
35. Dalmau J, Gleichman AJ, Hughes EG, Rossi JE, Peng X, Lai M, et al. Anti-NMDA-receptor encephalitis: case series and analysis of the effects of antibodies. *Lancet Neurol*. 2008;7(12):1091–8.
36. Nosadini M, Mohammad SS, Ramanathan S, Brilot F, Dale RC. Immune therapy in autoimmune encephalitis: a systematic review. *Expert Rev Neurother*. 2015;15(12):1391–419.
37. Chevdeville A, Honnorat J, Hampe CS, Desestret V. Neuronal central nervous system syndromes probably mediated by autoantibodies. *Eur J Neurosci*. 2016;43(12):1535–52.
38. Bergantini L, d'Alessandro M, Cameli P, Vietri L, Vagaggini C, Perrone A, Sestini P, Frediani B, Bargagli E. Effects of rituximab therapy on B cell differentiation and depletion. *Clin Rheumatol*. 2020;39(5):1415–21.
39. Dalakas MC. Invited article: inhibition of B cell functions: implications for neurology. *Neurology*. 2008;70(23):2252–60.
40. Dale RC, Brilot F, Duffy LV, Twilt M, Waldman AT, Narula S, et al. Utility and safety of rituximab in pediatric autoimmune and inflammatory CNS disease. *Neurology*. 2014;83(2):142–50.
41. Kaegi C, Wuest B, Schreiner J, Steiner UC, Vultaggio A, Matucci A, et al. Systematic review of safety and efficacy of rituximab in treating immune-mediated disorders. *Front Immunol*. 2019;10:1990. <https://doi.org/10.3389/fimmu.2019.01990>.
42. Chisari CG, Garlata E, Arena S, Toscano S, Luca M, Patti F. Rituximab for the treatment of multiple sclerosis: a review. *J Neurol*. 2022;269(1):159–83.
43. Zhao C, Li C, Duan FJ, Yan Q, Zhang Z, Du Y, Zhang W. Case report: repeated low-dose rituximab treatment is effective in relapsing neuro Behcet's disease. *Front Neurol*. 2021;12: 595984. <https://doi.org/10.3389/fneur.2021.595984>.
44. Du Y, Li C, Hao YF, Zhao C, Yan Q, Yao D, Li L, Zhang W. Individualized regimen of low-dose rituximab monotherapy for new-onset AChR-positive generalized myasthenia gravis. *J Neurol*. 2022. <https://doi.org/10.1007/s00415-022-11048-4>.
45. van Swieten JC, Koudstaal PJ, Visser MC, Schouten HJ, van Gijn J. Inter-observer agreement for the assessment of handicap in stroke patients. *Stroke*. 1988;19(5):604–7.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

