

RESEARCH

Open Access

Differential effects of interleukin-17 receptor signaling on innate and adaptive immunity during central nervous system bacterial infection

Debbie Vidlak¹ and Tammy Kielian^{1,2*}

Abstract

Although IL-17A (commonly referred to as IL-17) has been implicated in the pathogenesis of central nervous system (CNS) autoimmune disease, its role during CNS bacterial infections remains unclear. To evaluate the broader impact of IL-17 family members in the context of CNS infection, we utilized IL-17 receptor (IL-17R) knockout (KO) mice that lack the ability to respond to IL-17, IL-17F and IL-17E (IL-25). In this article, we demonstrate that IL-17R signaling regulates bacterial clearance as well as natural killer T (NKT) cell and gamma-delta ($\gamma\delta$) T cell infiltrates during *Staphylococcus aureus*-induced brain abscess formation. Specifically, when compared with wild-type (WT) animals, IL-17R KO mice exhibited elevated bacterial burdens at days 7 and 14 following *S. aureus* infection. Additionally, IL-17R KO animals displayed elevated neutrophil chemokine production, revealing the ability to compensate for the lack of IL-17R activity. Despite these differences, innate immune cell recruitment into brain abscesses was similar in IL-17R KO and WT mice, whereas IL-17R signaling exerted a greater influence on adaptive immune cell recruitment. In particular, $\gamma\delta$ T cell influx was increased in IL-17R KO mice at day 7 post-infection. In addition, NK1.1^{high} infiltrates were absent in brain abscesses of IL-17R KO animals and, surprisingly, were rarely detected in the livers of uninfected IL-17R KO mice. Although IL-17 is a key regulator of neutrophils in other infection models, our data implicate an important role for IL-17R signaling in regulating adaptive immunity during CNS bacterial infection.

Keywords: Brain abscess, IL-17R, Macrophages, $\gamma\delta$ T cells, Neutrophils, NKT cells

Introduction

Brain abscesses typically develop following parenchymal colonization with pyogenic bacteria, such as *Staphylococcus aureus* or *streptococcus* strains [1,2]. Characterized by an acute edematous response, *S. aureus* abscesses begin as localized areas of inflammation, evolving into suppurative lesions surrounded by a fibrotic capsule. Despite recent therapeutic advances, brain abscesses are still associated with significant morbidity and mortality [3]. In addition, long-term morbidity issues arise in patients recovering from these infections as a result of the extensive parenchymal damage typically associated with brain abscess formation, which can manifest as seizures, cognitive deficits, and/

or hemiparesis [3-5]. Because of the ubiquitous nature of bacteria and the continuous emergence of multi-drug resistant isolates, such as methicillin-resistant *S. aureus* (MRSA), these central nervous system (CNS) infections are likely to persist [6-8]. Therefore, a better understanding of the complex host-pathogen interactions that occur during brain abscess formation is essential for the development of novel therapies to treat these devastating infections.

The role of T helper 17 (Th17) cells in various inflammatory diseases has been a topic of intense investigation in recent years. Although Th17 cells have been implicated in the pathogenesis of autoimmune diseases [9-13], they have also been shown to provide protection against extracellular bacterial infections [14-16]. IL-17A (commonly referred to as IL-17) is the prototypic cytokine of the IL-17 family which includes six members, namely IL-17A, B, C, D, E and F [17-21]. In general, IL-17 plays an important role in regulating tissue inflammation through the

* Correspondence: tkielian@unmc.edu

¹Department of Pathology and Microbiology, University of Nebraska Medical Center, 985900 Nebraska Medical Center, Omaha, NE, 68198, USA

²Department of Pathology and Microbiology, University of Nebraska Medical Center, 985900 Nebraska Medical Center, Omaha, NE 68198-5900, USA

induction of proinflammatory cytokines (IL-6, TNF- α and IL-1 β) [22,23], chemokines (CXCL1, CXCL6, CXCL8 and CCL2) [24], and metalloproteinases [25] from various cell types, resulting in neutrophil recruitment [9,19,26]. To date, five IL-17 receptors (IL-17Rs) have been described (IL-17RA, -B, -C, -D and -E) [22,27-29], with mRNA ubiquitously expressed in a wide array of tissues (for example, lungs, kidney, liver, spleen and brain) [22,27] and multiple cell types (for example, macrophages, lymphocytes, fibroblasts and epithelial cells) [27]. The binding of IL-17 family members to their corresponding receptors triggers a signaling cascade that elicits cytokine and chemokine production. Because IL-17, IL-17E, and IL-17F all signal via the IL-17R [9,30,31], analysis of IL-17R knockout (KO) mice represents a way of investigating the broader impact of IL-17 family member action within the CNS. This was the rationale employed in the current study.

Among numerous proinflammatory mediators, IL-17 expression is elevated during brain abscess development [32]. Recent work from our laboratory has demonstrated that Th17 cells are critical for bacterial containment and regulation of innate immune cell infiltrates during the later stages of brain abscess formation [33,34]. This finding suggests that IL-17 may be a key determinant in regulating immune responses during CNS bacterial infection, but this remains to be determined as Th17 cells secrete other inflammatory cytokines that can affect the course of inflammation (for example, IL-22 and IL-27). Therefore, the current study was designed to evaluate the functional role of IL-17 family members in brain abscess pathogenesis by utilizing IL-17R KO mice. Our data revealed that IL-17R KO mice displayed higher bacterial burdens than wild-type (WT) animals, but that this had no effect on survival following CNS *S. aureus* infection. In accordance with the elevated expression of select inflammatory mediators (for example, CXCL2 and CXCL9), we also detected increased gamma-delta ($\gamma\delta$) T cell infiltrates in the brains of IL-17R KO mice, suggesting a potential compensatory mechanism in the absence of IL-17R signaling. Most notably, these studies describe an apparent natural killer T (NKT) cell deficiency in IL-17R KO animals, a novel finding that may offer insights into *S. aureus* CNS infection, as well as other peripheral models of infection and injury.

Materials and methods

Mice

IL-17RA KO mice (C57BL/6 background) were obtained from Amgen (Seattle, WA, USA) [9]. Age- and sex-matched C57BL/6 mice (Charles River Laboratories, Frederick, MD, USA, through a contract with National Cancer Institute) were used as WT controls. All animals were bred and housed in an AAALAC-accredited animal facility at the University of Nebraska Medical Center, provided with food and water *ad libitum*, and housed

under 12 h light/dark cycles. Brain abscess studies were performed with mice between 10 and 16 weeks of age.

Generation of experimental brain abscesses

Brain abscesses were induced by intracerebral injection of a MRSA USA300 strain encapsulated in agarose beads as previously described [35]. This isolate was recovered from an otherwise healthy individual who died from a brain abscess [36]. It is important to note that MRSA strains are uncommonly observed in community-acquired brain abscesses [36], whereas they are more prevalent in infections arising after trauma or neurosurgical procedures [37], and may differ in virulence compared to methicillin-sensitive *S. aureus* (MSSA). Briefly, mice were anesthetized with an intraperitoneal injection of 2.5% avertin. A 1 cm longitudinal incision was then made in the scalp to expose the underlying skull sutures and facilitate the identification of bregma. A rodent stereotaxic apparatus equipped with a Cunningham mouse adaptor (Stoelting, Kiel, WI, USA) was used to implant *S. aureus*-encapsulated beads into the striatum, using the following coordinates relative to bregma: +1.0 mm rostral, +2.0 mm lateral, and -3.0 mm deep from the surface of the brain. A burr hole was made and a 10 μ l Hamilton syringe fitted with a 26-gauge needle was used to slowly deliver 2 μ l of *S. aureus*-laden beads (7×10^3 - 1×10^4 colony forming units (CFU)) into the brain parenchyma. The needle remained in place for 2.5 minutes following injection to minimize bead efflux and potential leakage into the meninges. The skin incision was closed using surgical glue and animals were closely monitored over the course of each study for clinical indices of infection. The animal-use protocol, approved by the University of Nebraska Medical Center Animal Care and Use Committee, is in accord with the National Institutes of Health guidelines for the use of rodents.

Quantitation of viable bacteria from brain abscesses

To quantitate the numbers of viable *S. aureus* associated with brain abscesses *in vivo*, serial ten-fold dilutions of brain abscess homogenates were plated onto modified trypticase soy agar plates (Becton Dickinson, Sparks, MD, USA) supplemented with 5% defibrinated sheep blood (Hemostat Laboratories, Dixon, CA, USA). Titers were calculated by enumerating colonies and are expressed as CFU per gram of tissue.

Histological analysis of brain tissues

Immediately *ex vivo*, brains from IL-17R KO and WT mice were placed in a cryomold (Fisher Scientific, Fair Lawn, NJ, USA), embedded in Optimal Cutting Temperature (OCT) medium (Tissue-Tek, Torrance, CA, USA), and placed on dry ice until frozen. Cryostat sections (15 μ m) were mounted onto glass slides (Erie Scientific Co., Portsmouth, NH, USA) and subjected to H&E staining (Fisher

Scientific, Fair Lawn, NJ, USA). Images (20×) were collected using a digital slide scanner (Ventana Medical Systems, Tucson, AZ, USA) and the final images (1×) were prepared using Ventana Medical Image View Software.

Multi-analyte microbead array for detection of proinflammatory mediator production

To quantitate inflammatory mediator production in brain abscess homogenates, a mouse 19-plex microbead suspension array system was used according to the manufacturer's instructions (Millipore Corporation, Billerica, MA, USA). This customized array allows for the simultaneous detection of 19 individual inflammatory molecules in a single 50 µl sample, including IL-1α, IL-1β, TNF-α, IFN-γ, IL-6, IL-9, IL-10, IL-12 p40 and p70, IL-15, IL-17, CCL2, CCL3, CCL4, CCL5, CXCL1, CXCL2, CXCL9 and CXCL10. Results were analyzed using a Bio-Plex Workstation (Bio-Rad, Hercules, CA, USA) and adjusted based on the amount of total protein extracted from abscess tissues for normalization. The level of sensitivity for most analytes in the array was 3.2 pg/ml.

Quantitation of abscess-associated cells by fluorescence-activated cell sorting

To determine whether IL-17R signaling affected innate and/or adaptive immune cell influx into brain abscesses, cell populations were quantitated by fluorescence-activated cell sorting (FACS) as previously described [35,38,39]. Briefly, mice were manually perfused with isotonic PBS, pH 7.4, for approximately 2 minutes (at approximately 30 ml/minute) to eliminate leukocytes from the vasculature until the liver appeared blanched. Prior histological analysis had demonstrated the absence of leukocytes remaining adherent to the cerebral vascular endothelium and lack of perivascular cuffing in the Virchow Robin space of vascular-perfused mice in the brain abscess model (data not shown). Based on these observations, FACS analysis is an accurate representation of cells that have invaded the CNS parenchyma. Following vascular perfusion, the entire infected hemisphere was collected to recover abscess-associated cells, which ensured that equivalent tissue regions were obtained from both IL-17R KO and WT mice for downstream comparisons of leukocyte infiltrates. Tissues were minced in Hank's Balanced Salt Solution (HBSS; Hyclone Laboratories, Logan, UT, USA) supplemented with 10% Fetal Bovine Serum (FBS; Atlanta Biologicals, Lawrenceville, GA, USA) and filtered through a 70 µm nylon mesh cell strainer. At this point, an aliquot of tissue homogenate from each animal was collected to quantitate bacterial burdens. The resulting slurry was then digested for 30 minutes at 37 °C in HBSS supplemented with 0.2 mg/ml collagenase type I and 28 U/ml DNase I (both from Sigma-Aldrich, St Louis, MO, USA) to obtain a single-cell suspension. Following enzyme neutralization,

cells were layered onto a discontinuous Percoll gradient (1.03-1.088 g/ml) and centrifuged at 2,400 rpm for 20 minutes at room-temperature in a swinging bucket rotor. After centrifugation, myelin debris was carefully aspirated and the cell interface collected. Following extensive washes and incubation in Fc Block™ (BD Biosciences, San Diego, CA, USA), a panel of directly-conjugated antibodies was used for multi-color FACS to identify neutrophils (F4/80⁻, CD45⁺, Ly6G⁺), macrophages (F4/80⁺, CD45^{high}, Ly6G⁻), microglia (F4/80⁺, CD45^{low-intermediate}, Ly6G⁻), CD4 T cells (CD3⁺CD4⁺), CD8 T cells (CD8a⁺), NK cells (NKp46⁺, NK1.1⁺), NKT cells (CD3⁺, NKp46⁻, NK1.1⁺), and γδ T cells (γδTCR⁺). A recent study from our laboratory has established that the majority of CD4⁺ infiltrates during brain abscess development are Th1 and Th17 cells as shown by CD3⁺ co-expression [34]. Antibodies were purchased from the following vendors: F4/80-AlexaFluor488 (AbD Serotec, Raleigh, NC, USA); CD45-APC, Ly6G-PE, CD4-AlexaFluor700, CD8a-FITC, and NK1.1-APC (BD Biosciences); and NKp46-PE and γδTCR-PE-Cy5 (eBioscience, San Diego, CA, USA). Cells were analyzed using a BD LSRII (BD Biosciences) with compensation based on the staining of each individual fluorochrome alone and correction for autofluorescence with unstained cells. Controls included cells stained with isotype control antibodies to assess the degree of non-specific staining. Analysis was performed using BD FACSDiva™ software (BD Biosciences) with cells gated on the total leukocyte population. Results are presented as the absolute number of cells recovered from each brain, with normalization to adjust for the recovery of different cell numbers from the two mouse strains.

Statistics

Significant differences in bacterial titers, immune cell infiltrates, and proinflammatory mediator expression between IL-17R KO and WT mice at a particular time point were determined using a paired Student's *t*-test (SPSS Science, Chicago, IL, USA). A *P*-value of less than 0.05 was considered statistically significant.

Results



IL-17 receptor signaling is important for bacterial clearance during central nervous system *S. aureus* infection

Previous work from our laboratory has established significant IL-17 production and Th17 infiltrates during brain abscess development [32,34]. Based on these observations, it was expected that IL-17 family members would be an essential component in the developing immune response to *S. aureus* brain abscesses, since IL-17A and IL-17E are known to play a role in granulopoiesis, neutrophil recruitment, and proinflammatory cytokine production in response to extracellular bacterial infections [40,41]. To

investigate the functional importance of IL-17 signaling in the context of CNS infection, brain abscess pathogenesis was evaluated in IL-17R KO and WT mice. IL-17R signaling was found to be important in controlling bacterial clearance, as brain abscess tissues of IL-17R KO animals displayed significantly higher bacterial burdens at days 7 and 14 after infection compared with WT mice (Figure 1). Despite greater numbers of bacteria present in the brains of IL-17R KO mice, IL-17R signaling had no significant effect on survival in this model (Figure 2). Histological analysis of brain abscesses from IL-17R KO and WT animals did not reveal any significant alterations in lesion size between the two groups (Figure 3). In addition, no differences in edema formation were observed between IL-17R KO and WT mice at both the gross or histological levels, and abscess wet tissue weights between both groups were nearly identical (data not shown). Collectively, these findings suggest that edematous responses are not exaggerated in IL-17R KO animals following CNS bacterial infection.

Loss of IL-17 receptor signaling results in elevated proinflammatory mediator production in brain abscesses


The fact that IL-17 is known to induce neutrophil chemokine expression, in conjunction with our previous findings that IL-17-producing CD4⁺ T cells are the predominant T cell subset associated with brain abscesses [42], led us to predict that proinflammatory mediator production would be attenuated with the loss of IL-17R signaling. Surprisingly, the expression of several proinflammatory mediators was enhanced in IL-17R KO mice. While IL-17R KO animals predictably had significantly higher amounts of IL-17 in brain abscess homogenates (Figure 4A), several other mediators were also elevated



between 7 and 14 days after infection, including IL-1 α , IL-1 β , IL-6, CCL3 and CXCL1 (data not shown), as well as CXCL2 and CXCL9 (Figure 4B,D). These findings suggest that IL-17R signaling influences inflammatory mediator production on a more global scale, which may result from the inability of IL-17R KO mice to efficiently clear *S. aureus* from the CNS.

Innate immune cell recruitment into brain abscesses proceeds in an IL-17-independent manner

A primary role of IL-17 during tissue inflammation is neutrophil recruitment, mediated indirectly by the ability of IL-17 to induce neutrophil chemokine release [9,19,26]. However, the effect of IL-17R signaling on immune cell influx into the infected CNS is not known. To investigate this relationship, FACS analysis was performed on brain






Figure 4 Loss of IL-17 receptor signaling results in elevated proinflammatory mediator production. Supernatants were collected from brain abscess homogenates of IL-17 receptor (IL-17R) knockout (KO) and wild-type (WT) mice (n=5 to 8 per group) at days 7 and 14 after infection, whereupon IL-17 (a), CXCL2 (b), CCL2 (c) and CXCL9 (d) production was quantitated using multi-analyte bead arrays. Results are representative of two independent experiments and are presented as the average cytokine/chemokine concentration normalized to the total amount of protein (mean ± SD). * $P < 0.05$, ** $P < 0.01$, significant differences between IL-17R KO and WT mice.

abscess tissues from IL-17R KO and WT mice. Despite the important role that IL-17 plays in neutrophil recruitment in other model systems, it was not required during CNS *S. aureus* infection since IL-17R KO mice were as equally capable as WT animals of recruiting these cells to the site of infection (Figure 5A). The same trend was

observed with macrophage infiltrates (Figure 5B), and no noticeable differences in the absolute numbers of microglia were observed in these studies (data not shown). Collectively, these results indicate that IL-17R signaling has minimal effects on innate immune infiltrates during the course of CNS *S. aureus* infection.




Figure 5 IL-17 receptor signaling does not influence neutrophil or macrophage recruitment during central nervous system *S. aureus* infection. Abscess-associated cells from IL-17 receptor (IL-17R) knockout (KO) and wild-type (WT) mice (n=5 to 9 per group) were recovered at the indicated time points after infection and analyzed by flow cytometry to identify neutrophils and macrophages. Results are presented as the absolute cell number of each population recovered from a total of three independent experiments (mean ± SD).

IL-17 receptor loss differentially regulates T cell subset infiltration into the infected central nervous system

While previous studies from our laboratory have established the presence of IL-17-producing T cells during brain abscess development [32,34], we have yet to define how the action of IL-17R influences the accumulation of various T cell subsets. To address this issue, FACS analysis was performed on brain abscess tissues from IL-17R KO and WT animals. As previously demonstrated, CD4⁺ T cells were the most frequent T cell infiltrate, with fewer CD8⁺ T cells (Figure 6); however, no significant differences in the absolute numbers of CD4⁺ or CD8⁺ T cells were observed between IL-17R KO and WT mice (Figure 6). We expanded our analysis to include less frequent brain abscess T cell infiltrates, including NKT cells and $\gamma\delta$ T cells (Figure 7). Interestingly, NKT cells were significantly reduced in brain abscesses of IL-17R KO mice throughout infection (Figure 7A,B), whereas the opposite was true of $\gamma\delta$ T cells, which were significantly elevated compared with WT animals at day 7 post-infection (Figure 7A). In contrast, no significant differences in NK cell infiltrates were detected between groups (data not shown). These findings suggest that the loss of IL-17R signaling has the greatest effect on T cell populations that span innate-adaptive immunity (that is, NKT cells and $\gamma\delta$ T cells) in the context of CNS *S. aureus* infection.

IL-17 receptor knockout mice exhibit defects in natural killer T cell populations

Closer examination of our flow cytometry data revealed the presence of two distinct NK1.1⁺ populations in the

brain, namely NK1.1^{high} and NK1.1^{low} (Figure 8A,B). While there was little difference in NK1.1^{low} infiltrates between groups, the NK1.1^{high} population was significantly reduced in IL-17R KO mice, to the point of being difficult to detect in some experiments (Figure 8C,D). Based on these findings, we examined whether the failure to detect NK1.1^{high} infiltrates in brain abscesses resulted from a defect in recruitment or an inherent absence of these cells in the periphery. Therefore, the frequency of NK1.1^{high} cells was examined in the livers of uninfected IL-17R KO and WT mice as the majority of NK1.1^{high} cells reside in the liver [43]. Interestingly, NK1.1^{high} cells were rarely detected in the livers of non-manipulated IL-17R KO mice (Figure 9). To our knowledge, the requirement for IL-17R signaling in populating the liver with NK1.1^{high} cells has not yet been reported in the literature and could offer important insights into the relationship between IL-17R and T cell function in various animal models. However, since recent studies from our laboratory have demonstrated that NKT cells do not play a significant role in regulating immune responses during brain abscess development (Holley and Kielian, unpublished observations), this avenue was not pursued further.

Discussion

The role of IL-17 and its receptor have been well studied in recent years [20,44-46]. Numerous cell types are known to produce IL-17, including $\gamma\delta$, NKT and CD4⁺ T cells [47-49], resulting in robust inflammatory mediator production and subsequent neutrophil accumulation [22-24,49,50]. While IL-17 production is often associated with various autoimmune disorders [9-13], the cytokine is also known to exert protective effects during extracellular bacterial infections [14-16]. Our previous studies have demonstrated significant IL-17 production during brain abscess development [32-34], and Th17 cells represent a predominant CD4⁺ infiltrate. To understand the functional importance of IL-17 and related family members on brain abscess pathogenesis, we examined disease progression in IL-17R KO mice. Given its key role in neutrophil recruitment, and our previous findings that neutrophils are essential for survival during *S. aureus* abscess formation [51], we expected that loss of IL-17R signaling would result in a diminished capacity to control infection. To some extent this prediction was correct, as shown by the fact that IL-17R KO mice exhibited delayed bacterial clearance compared with WT animals. However, despite elevated bacterial burdens, survival rates were similar between IL-17R KO and WT animals. An explanation for the latter observation is that, despite the defect in IL-17R function, IL-17R KO mice were as effective as WT animals in recruiting innate immune cells into the infected brain. As established by previous studies in our laboratory [51], a correlation exists between the degree of neutrophil infiltration and bacterial






Figure 6 IL-17 receptor loss has no effect on T cell recruitment into brain abscesses. Abscess-associated cells from IL-17 receptor (IL-17R) knockout (KO) and wild-type (WT) mice (n = 5 to 9 per group) were recovered at the indicated time points after infection, and analyzed by flow cytometry to identify CD4⁺ and CD8⁺ T cells. Results are presented as the absolute cell number for each population recovered from a total of four independent experiments (mean \pm SD).



titers in the brain. Therefore, despite the elevated bacterial burdens observed in IL-17R KO mice, these animals were no more likely to succumb to infection than WT mice since they retained the ability to effectively recruit neutrophils into the brain. This suggests that an alternative signal (or signals) is capable of eliciting neutrophil chemokine expression and recruitment in the absence of IL-17R action. Indeed, many redundant mechanisms exist to ensure efficient pathogen recognition and clearance by the host immune system, with possible candidates including complement split products (for example, C3a, C5a) and bacterial components (for example, formylated peptides). Therefore, while IL-17R signaling is not essential for survival following CNS *S. aureus* challenge, it does appear to be important in controlling bacterial burdens, though the exact mechanism(s) of action remains to be determined.

Differentiation of naïve $CD4^+$ T cells into T helper or effector cells relies heavily on the inflammatory milieu. As we have previously demonstrated, Th17 infiltrates are

present in *S. aureus*-induced brain abscesses [33,34,42]. Based on the understanding that IL-17 plays a pivotal role in inducing and maintaining an effective immune response, as well as results from others utilizing IL-17R KO mice [9,52], we expected that a loss in IL-17R function would result in decreased proinflammatory mediator expression during CNS *S. aureus* infection. On the contrary, most mediators that we measured were elevated in brain abscesses of IL-17R KO animals. In particular, potent neutrophil chemoattractants, such as CXCL1 and CXCL2, were increased in IL-17R KO mice, possibly acting to compensate for the loss of IL-17R signaling. Evidence to support this possibility is provided by the fact that neutrophil infiltrates were similar in both IL-17R KO and WT animals. It is important to note that IL-17R KO mice have been shown to exhibit fewer circulating neutrophils in other studies [9,52]. This discrepancy may be explained by the fact that most reports using IL-17R KO mice have focused on peripheral sites of inflammation in



the spleen, liver and lungs, whereas we examined a unique tissue compartment in the CNS. Since IL-17RA is responsible for transducing signals emanating from IL-17, IL-17 F and IL-17E (IL-25) [20,29], we are not able to definitively assign the biological actions of IL-17R signaling to one specific IL-17 family member. Nonetheless, the approach to begin studying IL-17R KO animals was preferred since it represented a more global means of assessing IL-17 isoform involvement, whereupon the identification of specific family members can be assessed in future studies that are outside the scope of the current report.

While the early innate response is crucial to controlling bacterial burdens and recruiting effector cells into the brain during abscess formation, equally important is the developing adaptive immune response [34]. As we and others have previously shown [33,34,53], CD4⁺ T cells are

detected in brain abscesses as early as 3 days after infection and our recent study [34], using T cell adoptive transfers into TCR $\alpha\beta$ KO mice, revealed that both Th1 and Th17 cells are important for effective bacterial clearance during brain abscess development. To address whether Th1 or Th17 infiltrates were altered in the context of IL-17R loss, we performed intracellular cytokine staining for IL-17 and IFN- γ . No significant differences in the proportions of CD3⁺CD4⁺ Th1 or Th17 cells were observed in the brains of WT and IL-17R KO mice at day 7 following *S. aureus* infection (data not shown) when peak T cell infiltrates were apparent (Figure 6). As the absolute numbers of CD4⁺ T cells in the brains of IL-17R KO and WT mice were similar at day 7 post-infection, the increased IL-17 levels observed in KO mice are likely to be attributed to either cytokine accumulation based on receptor




Figure 9 NK1.1^{high} cells are rarely detected in the livers of IL-17 receptor knockout mice. Cells were collected from the livers of uninfected IL-17 receptor (IL-17R) knockout (KO) and wild-type (WT) mice (n = 3 to 6 per group) and analyzed by flow cytometry to identify NKT and gamma-delta ($\gamma\delta$) T cells. Fluorescence-activated cell sorting (FACS) contour plots are presented to demonstrate the different population densities and distributions between IL-17R WT and IL-17R KO mice. The total NK1.1⁺ population is indicated in red, whereas the total $\gamma\delta$ TCR⁺ population is indicated in green. Plots are representative of three independent experiments.

absence or, alternatively, the significantly larger population of $\gamma\delta$ T cells infiltrating the CNS of IL-17R KO mice that can also produce IL-17.

Most notable was the effect of IL-17R loss on non-traditional T cell populations, specifically NKT cells and $\gamma\delta$ T cells, which are often considered transitional cells because of their ability to bridge innate and adaptive immunity [54,55]. Our studies found that $\gamma\delta$ T cell infiltrates were significantly increased in brain abscesses of IL-17R KO mice compared with WT animals at day 7 after infection, and remained elevated through day 14. Cell surface expression of Toll-like receptor (TLR)2 on $\gamma\delta$ T cells [56], coupled with their ability to rapidly produce proinflammatory cytokines such as IL-17 [57,58] and IFN- γ [59,60], makes $\gamma\delta$ T cells adept as early responders to bacterial infection. However, the role that $\gamma\delta$ T cells play during brain abscess development in the context of IL-17R loss remains uncertain.

NKT cells represent a unique lymphocyte population that expresses NK cell markers as well as a semi-invariant T cell receptor [43,61] and NKT cell infiltrates are detected during early brain abscess development [42]. Initial studies reported significantly decreased accumulation of NKT cells in brain abscesses of IL-17R KO mice compared with WT animals at both days 7 and 14 following CNS infection. Further analysis revealed the presence of two distinct NK1.1⁺ populations, namely NK1.1^{high} and NK1.1^{low}. The NK1.1^{low}-expressing population was found to co-express the $\gamma\delta$ TCR, indicating

that these cells were likely to be $\gamma\delta$ T cells, albeit only a fraction of the total $\gamma\delta$ T cell population, as the absolute numbers of NK1.1⁺, $\gamma\delta$ TCR⁺ cells remained relatively constant throughout infection, with little difference noted between IL-17R KO and WT mice. Additionally, both the NK1.1^{high} and NK1.1^{low} cells in the brain were found to be predominantly CD4⁻ (data not shown). As these preliminary studies were designed to examine fundamental differences in infiltrating immune cells in IL-17R KO versus WT mice, a more comprehensive analysis would be required to determine whether these NK1.1⁺ populations represent classical or invariant NKT cells. However, although NKT cells represent a sizable immune cell infiltrate during early brain abscess development, recent studies from our laboratory utilizing CD1d KO mice, which lack all NKT cell subsets, failed to reveal any distinct phenotypes during CNS infection (Holley and Kielian, unpublished observations).

An intriguing finding of the current study was that NK1.1^{high} cell infiltrates were essentially absent in brain abscesses of IL-17R KO mice. We looked at whether defective NK1.1^{high} cell recruitment into brain abscesses in IL-17R KO animals resulted from impaired trafficking into the CNS or if these cells were absent in the periphery. While NK1.1⁺ cells are broadly distributed in mice, they are most frequent in the liver [43,61,62]. To determine whether IL-17R KO mice demonstrated an inherent deficiency in NK1.1^{high} cells, we analyzed liver tissues from uninfected IL-17R KO and WT animals. Interestingly,

FACS analysis revealed few NK1.1^{high}-expressing cells in IL-17R KO mice under resting conditions. Unlike the CD4⁺ NK1.1^{high} infiltrate in brain abscesses of WT animals, NK1.1^{high} cells in the livers of WT mice were predominantly CD4⁺ (data not shown). Several studies have described the immense diversity in NKT cell phenotypes and functionality [63-66]. The differential expression of CD4 on NK1.1⁺ cells isolated from the brain and liver of mice suggests that these may represent two functionally and potentially developmentally distinct populations. To our knowledge, this is the first report describing a paucity of NK1.1^{high} cells in IL-17R KO mice; a finding that could offer valuable insights into their function during both CNS and systemic diseases.

In summary, we have described an important role for IL-17R signaling in controlling bacterial burdens during CNS *S. aureus* infection. Despite their defect in *S. aureus* clearance, IL-17R KO mice were no more likely to succumb to infection than WT animals. This could be attributed to increased inflammatory infiltrates (that is, $\gamma\delta$ T cells) in brain abscess of IL-17R KO mice, corresponding with an ability to control infection in an IL-17-independent manner. Finally, we describe for the first time an inherent rarity of NK1.1^{high}-expressing cells in uninfected IL-17R KO mice. From a broader perspective, this finding could have important implications, as IL-17R KO mice are commonly used to study both autoimmune and infectious diseases [67,68].

Abbreviations

CCL2: Macrophage chemoattractant protein-1/MCP-1; CCL3: Macrophage inflammatory protein-1 α /MIP-1 α ; CXCL1: Keratinocyte chemokine/KC; CXCL2: Macrophage inflammatory protein-2/MIP-2; CXCL9: Monokine induced by IFN- γ /MIG; CFU: Colony forming units; CNS: Central nervous system; FACS: Fluorescence-activated cell sorting; $\gamma\delta$: Gamma-delta; H&E: Hematoxylin and eosin; IFN: Interferon; IL: Interleukin; IL-17R: IL-17 receptor; KO: Knockout; MRSA: Methicillin-resistant *Staphylococcus aureus*; MSSA: Methicillin-sensitive *S. aureus*; NK: Natural killer; NKT: Natural killer T; PBS: Phosphate-buffered saline; TH17: T helper 17; TLR: Toll-like receptor; TNF: Tumor necrosis factor; WT: Wild-type.

Competing interests

The authors declare that they have no competing interests.

Acknowledgements

This work was supported by the NIH National Institute of Neurological Disorders and Stroke (NINDS) R01 NS040730 to T.K. The authors thank Monica Holley, Amy Aldrich, and Amanda Angle for excellent technical assistance, Dr Costi Sifri for the USA300 isolate, Dr Charles Kuszynski, Megan Michalak, and Victoria Smith in the UNMC Cell Analysis Facility for assistance with FACS analysis, the UNMC Tissue Sciences Facility for staining and imaging brain tissue sections, and Ms Kari Nelson for editorial review of the manuscript.

Authors' contributions

DV performed the experiments and data analysis, participated in study design, and helped to draft the manuscript. TK conceived the study, participated in study design and data interpretation, and helped draft and revise the manuscript. Both authors have read and approved the final version of the manuscript.

Received: 4 January 2012 Accepted: 15 June 2012

Published: 15 June 2012

References

1. Townsend GC, Scheld WM: Infections of the central nervous system. *Adv Intern Med* 1998, **43**:403-447.
2. Mathisen GE, Johnson JP: Brain abscesses. *Clin Infect Dis* 1997, **25**:763-779.
3. Greenberg BM: Central nervous system infections in the intensive care unit. *Semin Neurol* 2008, **28**:682-689.
4. Davis LE, Baldwin NG: Brain abscess. *Curr Treat Options Neurol* 1999, **1**:157-166.
5. Lu CH, Chang WN, Lui CC: Strategies for the management of bacterial brain abscess. *J Clin Neurosci* 2006, **13**:979-985.
6. Carpenter J, Stapleton S, Holliman R: Retrospective analysis of 49 cases of brain abscess and review of the literature. *Eur J Clin Microbiol Infect Dis* 2007, **26**:1-11.
7. Jones ME, Draghi DC, Karlowsky JA, Sahn DF, Bradley JS: Prevalence of antimicrobial resistance in bacteria isolated from central nervous system specimens as reported by U.S. hospital laboratories from 2000 to 2002. *Ann Clin Microbiol Antimicrob* 2004, **3**:3.
8. Prasad KN, Mishra AM, Gupta D, Husain N, Husain M, Gupta RK: Analysis of microbial etiology and mortality in patients with brain abscess. *J Infect* 2006, **53**:221-227.
9. Ye P, Rodriguez FH, Kanaly S, Stocking KL, Schurr J, Schwarzenberger P, Oliver P, Huang W, Zhang P, Zhang J, Shellito JE, Bagby GJ, Nelson S, Charrier K, Peschon JJ, Kolls JK: Requirement of interleukin 17 receptor signaling for lung CXC chemokine and granulocyte colony-stimulating factor expression, neutrophil recruitment, and host defense. *J Exp Med* 2001, **194**:519-527.
10. Fujino S, Andoh A, Bamba S, Ogawa A, Hata K, Araki Y, Bamba T, Fujiyama Y: Increased expression of interleukin-17 in inflammatory bowel disease. *Gut* 2003, **52**:65-70.
11. Katz Y, Nadviv O, Beer Y: Interleukin-17 enhances tumor necrosis factor α -induced synthesis of interleukin 1, 6, and 8 in skin and synovial fibroblasts: a possible role as a "fine-tuning cytokine" in inflammation processes. *Arthritis Rheum* 2001, **44**:2176-2184.
12. Ziolkowska M, Koc A, Luszczkiewicz G, Ksiezopolska-Pietrzak K, Klimczak E, Chwalinska-Sadowska H, Maslinski W: High levels of IL-17 in rheumatoid arthritis patients: IL-15 triggers in vitro IL-17 production via cyclosporin A-sensitive mechanism. *J Immunol* 2000, **164**:2832-2838.
13. Lock C, Hermans G, Pedotti R, Brendolan A, Schadt E, Garren H, Langer-Gould A, Strober S, Cannella B, Allard J, Klonowski P, Austin A, Lad N, Kaminski N, Galli SJ, Oksenberg JR, Raine CS, Heller R, Steinman L: Gene-microarray analysis of multiple sclerosis lesions yields new targets validated in autoimmune encephalomyelitis. *Nat Med* 2002, **8**:500-508.
14. Higgins SC, Jarnicki AG, Lavelle EC, Mills KH: TLR4 mediates vaccine-induced protective cellular immunity to *Bordetella pertussis*: role of IL-17-producing T cells. *J Immunol* 2006, **177**:7980-7989.
15. Khader SA, Rickel EA: IL-23 and IL-17 in the establishment of protective pulmonary CD4⁺ T cell responses after vaccination and during *Mycobacterium tuberculosis* challenge. *Nat Immunol* 2007, **8**:369-377.
16. Yu JJ, Ruddy MJ, Wong GC, Sfintescu C, Baker PJ, Smith JB, Evans RT, Gaffen SL: An essential role for IL-17 in preventing pathogen-initiated bone destruction: recruitment of neutrophils to inflamed bone requires IL-17 receptor-dependent signals. *Blood* 2007, **109**:3794-3802.
17. Li H, Chen J, Huang A, Stinson J, Heldens S, Foster J, Dowd P, Gurney AL, Wood WI: Cloning and characterization of IL-17B and IL-17 C, two new members of the IL-17 cytokine family. *Proc Natl Acad Sci U S A* 2000, **97**:773-778.
18. Starnes T, Robertson MJ, Sledge G, Kelich S, Nakshatri H, Broxmeyer HE, Hromas R: Cutting edge: IL-17 F, a novel cytokine selectively expressed in activated T cells and monocytes, regulates angiogenesis and endothelial cell cytokine production. *J Immunol* 2001, **167**:4137-4140.
19. Hurst SD, Muchamuel T, Gorman DM, Gilbert JM, Clifford T, Kwan S, Menon S, Seymour B, Jackson C, Kung TT, Brieland JK, Zurawski SM, Chapman RW, Zurawski G, Coffman RL: New IL-17 family members promote Th1 or Th2 responses in the lung: in vivo function of the novel cytokine IL-25. *J Immunol* 2002, **169**:443-453.
20. Lee J, Ho WH, Maruoka M, Corpuz RT, Baldwin DT, Foster JS, Goddard AD, Yansura DG, Vandlen RL, Wood WI, Gurney AL: IL-17E, a novel proinflammatory ligand for the IL-17 receptor homolog IL-17Rh1. *J Biol Chem* 2001, **276**:1660-1664.

21. Fort MM, Cheung J, Yen D, Li J, Zurawski SM, Lo S, Menon S, Clifford T, Hunte B, Lesley R, Muchamuel T, Hurst SD, Zurawski G, Leach MW, Gorman DM, Rennick DM: **IL-25 induces IL-4, IL-5, and IL-13 and Th2-associated pathologies in vivo.** *Immunity* 2001, **15**:985–995.
22. Yao Z, Fanslow WC, Saelin MF, Rousseau AM, Painter SL, Comeau MR, Cohen JJ, Spriggs MK: **Herpesvirus Saimiri encodes a new cytokine, IL-17, which binds to a novel cytokine receptor.** *Immunity* 1995, **3**:811–821.
23. Jovanovic DV, Di Battista JA, Martel-Pelletier J, Jolicoeur FC, He Y, Zhang M, Mineau F, Pelletier JP: **IL-17 stimulates the production and expression of proinflammatory cytokines, IL-beta and TNF-alpha, by human macrophages.** *J Immunol* 1998, **160**:3513–3521.
24. Fossiez F, Djossou O, Chomarat P, Flores-Romo L, Ait-Yahia S, Maat C, Pin JJ, Garrone P, Garcia E, Saeland S, Blanchard D, Gaillard C, Das Mahapatra B, Rouvier E, Golstein P, Banchereau J, Lebecque S: **T cell interleukin-17 induces stromal cells to produce proinflammatory and hematopoietic cytokines.** *J Exp Med* 1996, **183**:2593–2603.
25. Park H, Li Z, Yang XO, Chang SH, Nurieva R, Wang YH, Wang Y, Hood L, Zhu Z, Tian Q, Dong C: **A distinct lineage of CD4 T cells regulates tissue inflammation by producing interleukin 17.** *Nat Immunol* 2005, **6**:1133–1141.
26. Ferretti S, Bonneau O, Dubois GR, Jones CE, Trifilieff A: **IL-17, produced by lymphocytes and neutrophils, is necessary for lipopolysaccharide-induced airway neutrophilia: IL-15 as a possible trigger.** *J Immunol* 2003, **170**:2106–2112.
27. Moseley TA, Haudenschild DR, Rose L, Reddi AH: **Interleukin-17 family and IL-17 receptors.** *Cytokine Growth Factor Rev* 2003, **14**:155–174.
28. Haudenschild D, Moseley T, Rose L, Reddi AH: **Soluble and transmembrane isoforms of novel interleukin-17 receptor-like protein by RNA splicing and expression in prostate cancer.** *J Biol Chem* 2002, **277**:4309–4316.
29. Clark HF, Gurney AL, Abaya E, Baker K, Baldwin D, Brush J, Chen J, Chow B, Chui C, Crowley C, Currell B, Deuel B, Dowd P, Eaton D, Foster J, Grimaldi C, Gu Q, Hass PE, Heldens S, Huang A, Kim HS, Klimowski L, Jin Y, Johnson S, Lee J, Lewis L, Liao D, Mark M, Robbie E, Sanchez C, et al: **The secreted protein discovery initiative (SPDI), a large-scale effort to identify novel human secreted and transmembrane proteins: a bioinformatics assessment.** *Genome Res* 2003, **13**:2265–2270.
30. Rickel EA, Siegel LA, Yoon BR, Rottman JB, Kugler DG, Swart DA, Anders PM, Tocker JE, Comeau MR, Budelsky AL: **Identification of functional roles for both IL-17RB and IL-17RA in mediating IL-25-induced activities.** *J Immunol* 2008, **181**:4299–4310.
31. Gaffen SL: **Structure and signalling in the IL-17 receptor family.** *Nat Rev Immunol* 2009, **9**:556–567.
32. Kielian T, Haney A, Mayes PM, Garg S, Esen N: **Toll-like receptor 2 modulates the proinflammatory milieu in Staphylococcus aureus-induced brain abscess.** *Infect Immun* 2005, **73**:7428–7435.
33. Nichols JR, Aldrich A, Mariani MM, Vidlak D, Esen N, Kielian T: **TLR2 deficiency leads to increased Th17 infiltrates in experimental brain abscesses.** *J Immunol* 2009, **182**:7119–7130.
34. Holley M, Kielian T: **Th1 and Th17 cells regulate innate immune responses and bacterial clearance during central nervous system infection.** *J Immunol*, . Manuscript in review.
35. Kielian T, Phulwani NK, Esen N, Syed MM, Haney AC, McCastlain K, Johnson J: **MyD88-dependent signals are essential for the host immune response in experimental brain abscess.** *J Immunol* 2007, **178**:4528–4537.
36. Sifri CD, Park J, Helm GA, Stempner ME, Shukla SK: **Fatal brain abscess due to community-associated methicillin-resistant Staphylococcus aureus strain USA300.** *Clin Infect Dis* 2007, **45**:e113–e117.
37. Schliamser SE, Backman K, Norrby SR: **Intracranial abscesses in adults: an analysis of 54 consecutive cases.** *Scand J Infect Dis* 1988, **20**:1–9.
38. Carson MJ, Reilly CR, Sutcliffe JG, Lo D: **Mature microglia resemble immature antigen-presenting cells.** *Glia* 1998, **22**:72–85.
39. Ford AL, Goodsall AL, Hickey WF, Sedgwick JD: **Normal adult ramified microglia separated from other central nervous system macrophages by flow cytometric sorting. Phenotypic differences defined and direct ex vivo antigen presentation to myelin basic protein-reactive CD4+ T cells compared.** *J Immunol* 1995, **154**:4309–4321.
40. Ouyang W, Kolls JK, Zheng Y: **The biological functions of T helper 17 cell effector cytokines in inflammation.** *Immunity* 2008, **28**:454–467.
41. Kolls JK, Linden A: **Interleukin-17 family members and inflammation.** *Immunity* 2004, **21**:467–476.
42. Vidlak D, Mariani MM, Aldrich A, Liu S, Kielian T: **Roles of Toll-like receptor 2 (TLR2) and superantigens on adaptive immune responses during CNS staphylococcal infection.** *Brain Behav Immun* 2011, **25**:905–914.
43. Godfrey DI, Hammond KJ, Poulton LD, Smyth MJ, Baxter AG: **NKT cells: facts, functions and fallacies.** *Immunity Today* 2000, **21**:573–583.
44. Lubberts E, Schwarzenberger P, Huang W, Schurr JR, Peschon JJ, van den Berg WB, Kolls JK: **Requirement of IL-17 receptor signaling in radiation-resistant cells in the joint for full progression of destructive synovitis.** *J Immunol* 2005, **175**:3360–3368.
45. McAllister F, Henry A, Kreindler JL, Dubin PJ, Ulrich L, Steele C, FINDER JD, Pilewski JM, Carreno BM, Goldman SJ, Pirhonen J, Kolls JK: **Role of IL-17A, IL-17 F, and the IL-17 receptor in regulating growth-related oncogene-alpha and granulocyte colony-stimulating factor in bronchial epithelium: implications for airway inflammation in cystic fibrosis.** *J Immunol* 2005, **175**:404–412.
46. Nagata T, McKinley L, Peschon JJ, Alcorn JF, Aujla SJ, Kolls JK: **Requirement of IL-17RA in Con A induced hepatitis and negative regulation of IL-17 production in mouse T cells.** *J Immunol* 2008, **181**:7473–7479.
47. Sutton CE, Lalor SJ, Sweeney CM, Brereton CF, Lavelle EC, Mills KH: **Interleukin-1 and IL-23 induce innate IL-17 production from gammadelta T cells, amplifying Th17 responses and autoimmunity.** *Immunity* 2009, **31**:331–341.
48. Michel ML, Mendes-da-Cruz D, Keller AC, Lochner M, Schneider E, Dy M, Eberl G, Leite-de-Moraes MC: **Critical role of ROR-gamma in a new thymic pathway leading to IL-17-producing invariant NKT cell differentiation.** *Proc Natl Acad Sci U S A* 2008, **105**:19845–19850.
49. Yao Z, Painter SL, Fanslow WC, Ulrich D, Macduff BM, Spriggs MK, Armitage RJ: **Human IL-17: a novel cytokine derived from T cells.** *J Immunol* 1995, **155**:5483–5486.
50. Laan M, Cui ZH, Hoshino H, Lotvall J, Sjostrand M, Gruenert DC, Skoogh BE, Linden A: **Neutrophil recruitment by human IL-17 via C-X-C chemokine release in the airways.** *J Immunol* 1999, **162**:2347–2352.
51. Kielian T, Barry B, Hickey WF: **CXC chemokine receptor-2 ligands are required for neutrophil-mediated host defense in experimental brain abscesses.** *J Immunol* 2001, **166**:4634–4643.
52. Smith E, Stark MA, Zarbock A, Burcin TL, Bruce AC, Vaswani D, Foley P, Ley K: **IL-17A inhibits the expansion of IL-17A-producing T cells in mice through "short-loop" inhibition via IL-17 receptor.** *J Immunol* 2008, **181**:1357–1364.
53. Stenzel W, Soltek S, Sanchez-Ruiz M, Akira S, Miletic H, Schluter D, Deckert M: **Both TLR2 and TLR4 are required for the effective immune response in Staphylococcus aureus-induced experimental murine brain abscess.** *Am J Pathol* 2008, **172**:132–145.
54. Tupin E, Kinjo Y, Kronenberg M: **The unique role of natural killer T cells in the response to microorganisms.** *Nat Rev Microbiol* 2007, **5**:405–417.
55. Born WK, Yin Z, Hahn YS, Sun D, O'Brien RL: **Analysis of gamma delta T cell functions in the mouse.** *J Immunol* 2010, **184**:4055–4061.
56. Hayday AC: **Gammadelta T cells and the lymphoid stress-surveillance response.** *Immunity* 2009, **31**:184–196.
57. Lockhart E, Green AM, Flynn JL: **IL-17 production is dominated by gammadelta T cells rather than CD4 T cells during Mycobacterium tuberculosis infection.** *J Immunol* 2006, **177**:4662–4669.
58. Shibata K, Yamada H, Hara H, Kishihara K, Yoshikai Y: **Resident Vδ1+ γδ T cells control early infiltration of neutrophils after Escherichia coli infection via IL-17 production.** *J Immunol* 2007, **178**:4466–4472.
59. Lochner M, Peduto L, Cherrier M, Sawa S, Langa F, Varona R, Riethmacher D, Si-Tahar M, Di Santo JP, Eberl G: **In vivo equilibrium of proinflammatory IL-17+ and regulatory IL-10+ Foxp3+ RORgamma t+ T cells.** *J Exp Med* 2008, **205**:1381–1393.
60. Ivanov II, McKenzie BS, Zhou L, Tadokoro CE, Lepelletier JJ, Cua DJ, Littman DR: **The orphan nuclear receptor RORgamma directs the differentiation program of proinflammatory IL-17+ T helper cells.** *Cell* 2006, **126**:1121–1133.
61. Bendelac A, Rivera MN, Park SH, Roark JH: **Mouse CD1-specific NK1 T cells: development, specificity, and function.** *Annu Rev Immunol* 1997, **15**:535–562.
62. MacDonald HR: **NK1.1+ T cell receptor-alpha/beta+ cells: new clues to their origin, specificity, and function.** *J Exp Med* 1995, **182**:633–638.
63. Eberl G, Lees R, Smiley ST, Taniguchi M, Grusby MJ, MacDonald HR: **Tissue-specific segregation of CD1d-dependent and CD1d-independent NK T cells.** *J Immunol* 1999, **162**:6410–6419.
64. Hammond KJ, Pelikan SB, Crowe NY, Randle-Barrett E, Nakayama T, Taniguchi M, Smyth MJ, van Driel IR, Scollay R, Baxter AG, Godfrey DI: **NKT cells are phenotypically and functionally diverse.** *Eur J Immunol* 1999, **29**:3768–3781.

65. Zeng D, Gazit G, Dejbakhsh-Jones S, Balk SP, Snapper S, Taniguchi M, Strober S: **Heterogeneity of NK1.1+ T cells in the bone marrow: divergence from the thymus.** *J Immunol* 1999, **163**:5338–5345.
66. Apostolou I, Cumano A, Gachelin G, Kourilsky P: **Evidence for two subgroups of CD4-CD8- NKT cells with distinct TCR alpha beta repertoires and differential distribution in lymphoid tissues.** *J Immunol* 2000, **165**:2481–2490.
67. Gonzalez-Garcia I, Zhao Y, Ju S, Gu Q, Liu L, Kolls JK, Lu B: **IL-17 signaling-independent central nervous system autoimmunity is negatively regulated by TGF-beta.** *J Immunol* 2009, **182**:2665–2671.
68. Zhang Z, Zheng M, Bindas J, Schwarzenberger P, Kolls JK: **Critical role of IL-17 receptor signaling in acute TNBS-induced colitis.** *Inflamm Bowel Dis* 2006, **12**:382–388.

doi:10.1186/1742-2094-9-128

Cite this article as: Vidlak and Kielian: Differential effects of interleukin-17 receptor signaling on innate and adaptive immunity during central nervous system bacterial infection. *Journal of Neuroinflammation* 2012 **9**:128.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

 **BioMed** Central