

CORRECTION

Open Access



Correction to: the disruption of mitochondrial axonal transport is an early event in neuroinflammation

Oihana Errea¹, Beatriz Moreno¹, Alba Gonzalez-Franquesa², Pablo M. Garcia-Roves^{2,3,4} and Pablo Villoslada^{1,5*}

Correction

After publication of the article [1], it has been brought to our attention that the full funding acknowledgement is missing from the original article. It should also include the following –

“This work was supported by the Instituto de Salud Carlos III with FEDER funds (Otra forma de hacer Europa) from the European Commission (FIS: PI12/01823)”.

Author details

¹Center of Neuroimmunology, Institut d'Investigacions Biomèdiques August Pi Sunyer (IDIBAPS), Cellex Building, Laboratory 3A, Casanova 145, 08036 Barcelona, Spain. ²Diabetes and Obesity Research Laboratory, Institut d'Investigacions Biomèdiques August Pi Sunyer (IDIBAPS), 08036 Barcelona, Spain. ³Spanish Biomedical Research Center in Diabetes and associated disorders (CIBERDEM), University of Barcelona, 08907 Barcelona, Spain. ⁴Department of Physiological Sciences II, University of Barcelona, 08907 Barcelona, Spain. ⁵University of California, San Francisco, USA.

Published online: 23 October 2017

Reference

1. Errea O, Moreno B, Gonzalez-Franquesa A, Garcia-Roves P, Villoslada P. The disruption of mitochondrial axonal transport is an early event in neuroinflammation. *J Neuroinflammation*. 2015;12(1):152. doi:10.1186/s12974-015-0375-8.

* Correspondence: pvillosl@clinic.cat

¹Center of Neuroimmunology, Institut d'Investigacions Biomèdiques August Pi Sunyer (IDIBAPS), Cellex Building, Laboratory 3A, Casanova 145, 08036 Barcelona, Spain

⁵University of California, San Francisco, USA

